

**METHODS FOR PREDICTING THE BIOLOGICAL,  
CHEMICAL, AND PHYSICAL PROPERTIES OF  
MOLECULES FROM THEIR SPECTRAL PROPERTIES**

5

**FIELD OF THE INVENTION**

The invention relates to methods for predicting the biological, chemical, and physical properties of molecules from their spectral properties.

10

**BACKGROUND**

The number of known chemical compounds is vast, and the number is increasing constantly because methods for isolating and synthesizing molecules continue to improve. For instance, chemists are now able to employ the techniques of combinatorial chemistry to synthesize thousands of different chemical compounds at one time using a mixture of only a few interchangeable chemical building blocks. Furthermore, chemists are now able to use combinatorial computer models to generate large numbers of chemical structures that may in theory be synthesized.

While there are many chemical compounds, only a relative few of those compounds may exhibit a particular desirable property, such as pharmaceutical activity. Random testing of known compounds to find those that show pharmaceutical activity is very expensive and time-consuming. Similarly, there is a need to screen compounds for toxicity, so that rational decisions can be made regarding the use and regulation of compounds that have toxic potential. At present, only a fraction of known compounds have been thoroughly tested for their toxicological and potential therapeutic properties. To address this problem, scientists have developed methods which attempt to predict which compounds are likely to exhibit a particular property.

Methods for predicting the properties of chemical compounds are generally based upon the related observations that the structure of a compound is related to its biological, chemical, and physical properties, and that compounds of similar structure exhibit similar properties. These observations have been used to search for new compounds exhibiting a particular property. For instance, a benzene ring

is present in both acetaminophen and salicylamide, both of which are analgesics. Although incorporating a benzene ring into a new molecule increases the likelihood that it too will exhibit analgesic activity, this deduction only narrows the compounds to be tested. This approach is still basically one of trial and error,  
5 because many compounds with a benzene ring are not analgesics. Moreover, analgesics without a benzene ring will be missed in the search.

Quantitative structure-property relationships and quantitative structure-activity relationships (collectively QSAR) are attempts to quantify the observed relationships between the structure of chemical compounds and the extent to which  
10 those compounds exhibit certain properties. For instance, a QSAR might attempt to quantify how the analgesic activity of known analgesics that contain a benzene ring (such as acetaminophen and salicylamide) depends upon the number and identity of substituents on their benzene rings. Once established, such a QSAR could be used to predict the analgesic activity of other compounds that contain  
15 benzene rings, and identify those compounds that warrant further investigation as analgesics based on their predicted analgesic activity.

In the terminology associated with the QSAR method, the property for which a prediction is sought, such as analgesic activity, is termed the "endpoint." In general the endpoint may be any measurable biological, chemical or physical  
20 property.

To establish a QSAR, endpoint values are obtained for a set of compounds and a correlation is then sought between the endpoint values and some measure(s) of structure available for each of the compounds. The measures used to describe or reflect the structure of the compounds for which a correlation is sought are  
25 termed structure descriptors. Structure descriptors may be defined directly with reference to the known structures of the compounds, or may indirectly reflect the structure through a property of the molecule that is sensitive to changes in structure. For example, an investigator might try to correlate the analgesic activity of compounds that contain a benzene ring with either a direct measure of the  
30 structure, such as the number of hydroxyl groups attached to the benzene ring, or an indirect measure of the structure of the compounds, like water solubility. If the

direct measure is chosen, the attempted correlation could only include those compounds with hydroxyl groups on the benzene ring, while the indirect measure is more general and could be used to include all benzene ring containing compounds in the attempted correlation.

5           The endpoint data and the structure descriptor(s) for the set of compounds that are chosen to establish a QSAR are termed the training set. In general, the reliability of a QSAR increases as the number of compounds in the training set increases. If possible, the training set desirably includes compounds that exhibit a wide range of endpoint values and possess diverse structures.

10           If the endpoint and structure descriptor(s) are sufficiently correlated, a mathematical or graphical representation may be obtained. For example, the growth inhibition of certain gram negative bacteria by aromatic amines is correlated in a linear fashion with the logarithm of the octanol-water partition coefficient, and the correlation indicates that as the aromatic amines become more  
15 hydrophobic they are more likely to inhibit the growth of these bacteria (Hansch and Leo, *Exploring QSAR: Fundamentals and Applications in Chemistry and Biology*, American Chemical Society, 1995, p. 416).

          The QSAR representation may be used to predict endpoint values for other compounds from their structure descriptor(s), and the reliability of a QSAR may  
20 be tested using a validation set of data. A validation set includes structural and endpoint data for compounds that were not part of the training set. The validation set desirably exhibits a diversity of endpoint values and structures that is commensurate with the training set. The QSAR is tested by how reliably it predicts endpoint data for the validation set compounds from the validation set  
25 structural data. For example,  $\alpha$ -naphthylamine, an aromatic amine, is about two times less toxic than predicted by its octanol-water partition coefficient, indicating that the simple linear relationship described above is not always reliable and that some specific interaction is responsible for its behavior.

          The Hammet equation is another example of a simple QSAR that relates a  
30 single structure descriptor (in this case, derived directly from the structure of the compounds) to an endpoint (in this case, the equilibrium or rate constant for a

particular type of reaction). The Hammett equation is a linear equation of the form:

$$\ln K_x = \rho\sigma + \ln K_H$$

- 5 The electronic parameter ( $\sigma$ ) is a measure of the ability of a group of atoms (typically a substituent at a particular structural position) to donate (negative value) or withdraw (positive value) electron density to or from the reaction center of the molecule to which it is attached. The slope parameter ( $\rho$ ) is a measure of the sensitivity of the reaction to the withdrawal or release of electron density, and is
- 10 constant for a particular type of reaction.  $K_x$  and  $K_H$  are, respectively, the equilibrium constant for reaction of the substituted molecule and the equilibrium constant for reaction of the unsubstituted parent molecule. The reaction rate constants,  $k_x$  and  $k_H$ , may replace the equilibrium constants. A plot of  $\ln K_x$  (or  $\ln k_x$ ) versus  $\sigma$  is often linear and may be used, for example, to predict the
- 15 equilibrium constant (or rate constant) for other structurally similar compounds from only the  $\sigma$  values of their substituents.

The Hammett parameter  $\sigma$  is an example of a structure descriptor that is derived from experimental data for compounds of known structure. As described above,  $\sigma$  is a measure of the electronic properties of a substituent. Specifically  $\sigma$

20 measures the ability of a substituent at a particular position to donate or withdraw electron density and may be defined according to the following equation:

$$\sigma = \log K_x - \log K_H$$

- 25 where  $K_x$  and  $K_H$  are, respectively, the acid ionization constants for an aromatically substituted benzoic acid derivative and for unsubstituted benzoic acid. The parameter  $\sigma$  for amino group substitution in a position para to the reaction center is determined, for example, from the measured acid ionization constants of *p*-aminobenzoic acid and benzoic acid using the above equation. Although
- 30 depending on the position of substitution, electron-withdrawing substituents generally tend to stabilize the anion formed when benzoic acid ionizes, making  $K_x$  larger than  $K_H$  and  $\sigma$  positive. Conversely, electron-donating substituents tend to

destabilize the anion and generally have negative  $\sigma$  values that are also dependent upon the position of substitution. Acid dissociation data for benzoic acid derivatives with various substituents in ortho, meta or para positions have been used to generate  $\sigma$  values for many substituents in these structural positions.

- 5 Unfortunately, the  $\sigma$  values derived for substituents in this manner are typically not valid for multiply substituted molecules because  $\sigma$  values may not be additive.

Due to the inability of  $\sigma$  values to accurately reflect the cumulative electronic effect of multiple substituents, alternative structure descriptors have been investigated. For example, Ijzerman et al. derived structure descriptors from the assigned  $^{13}\text{C}$  NMR shifts of the aromatic carbons in a family of N-*tert*-butylphenylethanolamines (Ijzerman et al., *J. Med. Chem.*, **29**: 549-554, 1986). Ijzerman et al. subtracted the value of the chemical shift for the carbon atoms in benzene (128.5ppm) from the assigned chemical shift values for each of the carbons in the benzene ring of each of the substituted N-*tert*-butylphenylethanolamines to yield sets of structure descriptors for the compounds. A correlation was found between certain linear combinations of these structure descriptors and the  $\beta_2$ -adrenoreceptor intrinsic activity of the substituted N-*tert*-butylphenylethanolamines. Since these structure descriptors are defined in a manner similar to the Hammett  $\sigma$  parameter, they necessarily rely upon knowledge of structure beforehand. Also, the scope of usefulness for this type of structure descriptor is limited to the compounds containing a benzene ring. Furthermore, these structure descriptors ignore the aliphatic carbon atoms of the molecules, and their contribution to the intrinsic activity (a valid assumption only when comparing the intrinsic activities of compounds that differ minimally in structure).

25 Additionally, the correlations of Ijzerman et al. have been questioned because the structure descriptors they derived from the chemical shifts of the aromatic carbons in these compounds are themselves highly correlated, and their use leads to the unrealistic result that substitution at one ortho position of the benzene ring affects intrinsic activity in a manner opposite to substitution at the other ortho position (see, Cramer et al., *Quant. Struct.-Act. Relat.*, **7**: 18-25, 1988).

30

A similar attempt to utilize spectroscopically derived parameters as structure descriptors was made by Nishikawa and Tori (Nishikawa and Tori, *J. Med. Chem.*, **27**: 1657-1663, 1984). The reactivity of the  $\beta$ -lactam ring of 3-substituted and 3-methylene substituted cephalosporins toward alkaline hydrolysis showed a correlation with changes in the  $^{13}\text{C}$  NMR chemical shift of selected carbon atoms, and with changes in the infrared (IR) stretching frequency observed for the  $\beta$ -lactam carbon-oxygen bond. Selection of these measures was guided by knowledge of the structure of these compounds, and the role  $\beta$ -lactam reactivity plays in their antibiotic activity. Like the method of Ijzerman et al., the method of Nishikawa defines its structure descriptors in terms based upon a reference compound of known structure and utilizes spectral features assigned to particular atoms of known structures. Furthermore, the method, through its definition of the structure descriptors, is similarly restricted to narrow classes of compounds that differ only by simple substitutions.

Structure descriptors may be obtained theoretically or experimentally. Dipole moments and lowest unoccupied molecular orbital (LUMO) energies are examples of structure descriptors that may be obtained from theoretical quantum mechanical calculations on known structures. Experimental data correlated with a specific structural feature, common to a set of closely related compounds, may also be used to generate measures of structure (for example, the Hammett  $\sigma$  parameter). Bulk experimental measures of structure, such as partition coefficients (as a measure of polarity) and molar refractivities (as a measure of steric size) can also be utilized as structure descriptors. Structure descriptors based upon bulk physical properties have the advantage that they do not require structural knowledge beforehand, however such descriptors lack specificity. For example, compounds with vastly different biological activities may have very similar partition coefficients.

A particularly important type of biological QSAR uses, as an endpoint, the ability of one molecule to bind to another molecule. An example of such an endpoint would be the ability of a series of molecules to act as ligands for a regulatory protein, such as a hormone receptor. In recognition of the important

role played by three-dimensional structure, especially for biochemical reactions where molecular recognition is an important factor, the field of 3D-QSAR has emerged. The 3D-QSAR technique is exemplified by the Comparative Molecular Field Analysis (CoMFA) method of Cramer and Wold (U.S. Patent No.

5 5,025,388). The CoMFA method attempts to correlate the three-dimensional steric and electrostatic properties of a series of molecules with their relative endpoint values. The steric and electrostatic properties of a molecule are obtained from quantum mechanical or electrostatic calculations based upon known molecular structures and serve as structure descriptors. The calculations, in effect,  
10 map the electron density distribution around a molecule to create a 3-D picture of its steric and electrostatic fields (collectively, the molecular field). Those steric features (e.g., bulky substituents) and/or electrostatic properties (e.g., a strong molecular dipole) that are most important in determining endpoint values are revealed by comparing the molecular fields of the molecules in the training set to  
15 their endpoints. The advantage of calculated 3D-QSAR molecular field structure descriptors is that unlike structure descriptors referenced to a certain structural feature, molecular field structure descriptors enable the identification of structurally dissimilar molecules that have similar steric and electrostatic properties.

20 However, a particular problem associated with the CoMFA method and other 3D-QSAR techniques is that these methods generally require some assumptions about how molecules orient themselves relative to each other upon binding. Selecting the common alignment of a training set containing diverse structures may be problematic, leading to incorrect predictions of binding ability.  
25 Furthermore, QSAR based upon quantum mechanical or electrostatic potential calculations also suffers to some extent from the inaccuracy of the calculations themselves. These calculations are by nature approximate, and become less and less reliable as molecular size increases. In addition, the effects of solvation on the quantum mechanical properties of a molecule are often difficult to calculate and  
30 typically are ignored. Finally, these calculations are time-consuming and require knowledge of the molecular structures beforehand.

Isolation and structure elucidation of molecules is an expensive and time-consuming process that is desirably avoided, especially when screening large numbers of molecules (e.g., those generated in a combinatorial library).

Furthermore, many molecular structures are proprietary, so there may be  
5 reluctance on the part of investigators to share training set structural and endpoint data with competitors in a common effort to improve the reliability of predictive QSARs.

It would be advantageous to have QSAR methods that utilize structure descriptors which may be obtained without knowledge of or disclosure of  
10 molecular structures. Such structure descriptors could be more specific than bulk physical properties, and enable the QSAR methods to differentiate dissimilar molecular structures that exhibit similar bulk physical properties. Furthermore, the structure descriptors could avoid definition with reference to a particular structural element (e.g., the Hammett  $\sigma$  parameter), and therefore be useful for  
15 establishing QSARs for more structurally diverse sets of compounds. Additionally, such QSAR methods could eliminate the costly step of structure elucidation and obviate the need for isolation of the subject compounds.

QSAR methods that utilize structure descriptors that are inherently reflective of the steric and electrostatic properties of molecules and the effects of  
20 solvation thereon are also needed. Such methods would also desirably eliminate the necessity for assumptions regarding molecular orientation in relationship to intermolecular binding and obviate the need to rely upon approximate calculations.

#### SUMMARY OF THE DISCLOSURE

The present invention avoids some of the foregoing problems by providing  
25 a method for predicting a biological activity of a molecule, by obtaining spectral data (such as NMR data) for a test compound, and comparing the spectral data for a test compound to a pattern derived not exclusively from the assigned spectral data (such as NMR data) of a training set of compounds having known biological activity. Similarities between the pattern of spectral data associated with the  
30 biological activity of the training set compounds and the spectral data for the test compound are detected to determine whether the test compound is predicted to



exhibit the biological activity. The spectral data of the compound for which a prediction is sought need not first be correlated with corresponding structural features. Furthermore, the pattern of spectral data associated with the biological activity may be derived without first correlating the spectral data with

5 corresponding structural features. Training set patterns and similarities between the training set patterns and the test compound's spectral data are conveniently detected, in some embodiments, by segmenting the spectral data of the training set and test compounds into sub-spectral units (bins). These sub-spectral units, or bins, may be of a width corresponding to the digital resolution of the method used  
10 to generate the spectral data or greater.

For example, the biological activity of a test compound may be predicted by comparing the signals in bins of the training set spectral data that are found to be associated with a biological activity (such as strong estrogen receptor binding) to signals in corresponding bins of the spectral data of the test compound.

15 Numerous signals in the test compound's spectral data that fall within bins corresponding to strong estrogen receptor binding are an indication that the test compound possesses strong estrogen receptor binding. Some of the bins of the training set spectral data may contain signals more consistently associated with strong estrogen receptor binding, in which case the presence of signals in the  
20 corresponding bins for the test compound would be more heavily weighted in assigning a predicted biological activity to the test compound. Lesser numbers of signals in the test compound's spectral data that fall within bins corresponding to strong estrogen receptor binding indicate that the test compound possess only moderate or weak estrogen receptor binding.

25 The spectral data of the training compounds and the test compound may be just one type of spectral data (such as NMR, for example  $^{13}\text{C}$ -NMR), or more than one type of spectral data (such as a composite of two or more of NMR, mass spectral, infrared, ultraviolet-visible, fluorescence, or phosphorescence data). In particular embodiments, the spectral data is a composite of two or more of nuclear  
30 magnetic resonance spectroscopic (NMR) data, mass spectroscopic (MS) data,

infrared (IR) spectroscopic data, and ultraviolet-visible (UV-Vis) spectroscopic data.

In other embodiments, the spectral data of the training set compounds is segmented into sub-spectral units (bins), and scaled to normalize the importance of different signals (e.g., those from different types of spectra in a composite or those arising from signals of different inherent intensities) prior to pattern recognition. In particular embodiments, the scaling is auto-scaling. In yet other embodiments, the spectral data of the training set compounds is weighted prior to pattern recognition to emphasize those sub-spectral units (bins) that are most important for differentiating endpoint classes (such as strong versus weak estrogen receptor binding) of compounds in the training set. In more particular embodiments, the weighting is Fisher-weighting.

A pattern of spectral data associated with a biological activity can be advantageously derived from the training set spectral data using computer implemented pattern recognition techniques. Furthermore detection of similarities between the pattern derived from the training set spectral data and the spectral data exhibited by a test compound is also advantageously performed using computer implemented methods. In one such approach, the pattern of spectral data associated with a biological activity is derived for a training set of compounds by segmenting the spectral data and generating a set of canonical variate factors, one for each bin of the segmented spectral data. These canonical variate factors are used with the spectral data of a test compound to yield a prediction of the biological activity of the test compounds.

The methods of the present invention are advantageously computer implemented. In one such embodiment, the method is a computer implemented system for predicting biological activity of a test compound, in which input spectral data is received for a test compound, and for a set of training compounds having a known biological activity. The spectral pattern derived from the training set (derived, for example, using computer implemented pattern recognition programs) and the spectral pattern of the test compound are compared to determine whether the spectral patterns of the test compound match spectral patterns of the

training set associated with a biological activity. The spectral data for the test compound and the spectral data for the training set may conveniently be divided into substantially identical spectral bins, so that a signal within individual corresponding spectral bins is compared between the pattern derived from the  
5 training set and the test compound.

In this computer implemented system, the spectral patterns are obtained by inputting spectral data such as one or more of nuclear magnetic resonance data, mass spectral data, infrared data, ultraviolet-visible data, fluorescence data, and phosphorescence data. In particular embodiments, the spectral data is a composite  
10 of nuclear magnetic resonance data and mass spectral data. The spectral data of the training set is converted into principal components (PCs) and canonical variates (CVs). Peaks in particular bins of the canonical variates that are associated with a biological activity (such as high affinity to a hormone receptor) are identified, and the test compound is analyzed for the presence of one or more (and ideally many)  
15 corresponding peaks in its composite spectrum.

Another aspect of the invention is a method for predicting a biological, chemical, or physical property of molecules, by providing spectral data segmented into spectral sub-units, for a plurality of training compounds; inputting the segmented spectral data and endpoint data into a pattern-recognition program;  
20 training the pattern-recognition program with the segmented spectral data and endpoint data to establish a relationship between the spectral sub-units of the segmented spectral data and the endpoint; providing segmented spectral data for a test compound that is segmented into substantially the same spectral sub-units that were used for the training set, and comparing the relationship between the spectral  
25 sub-units of the segmented spectral data and the endpoint to the spectral sub-units of the test compound's segmented data to predict the endpoint of the test compound. In this method, the structures of the training compounds and the test compound are not necessarily known beforehand. In particular embodiments of the method, the segmented spectral data of the training set is auto-scaled and  
30 Fisher-weighted before training the pattern recognition program.

The spectral data is chosen from the group consisting of nuclear magnetic resonance data, mass spectral data, infrared data, UV-Vis data, fluorescence data, phosphorescence data, and composites thereof, for example  $^{13}\text{C}$  NMR data, EI MS data, and composites thereof. The endpoint may be a ligand-target molecule-binding affinity, such as estrogen-receptor binding affinity. Other examples of the endpoint are a measure of biodegradability; a measure of toxicity; participation in a metabolic pathway; a partition coefficient; a reaction rate; a quantum yield, a measure of phototoxicity, an equilibrium constant; and a site of reaction on a molecular structure. In a particular embodiment, the endpoint is the octanol/water partition coefficient.

Additional non-spectral structure descriptors may be utilized along with segmented spectral data to provide an expanded set of structure descriptors useful for establishing a predictive relationship for the endpoint. Examples of non-spectral structure descriptors that do not necessarily require structural knowledge beforehand include partition coefficients, solubilities, relative acidities, relative basicities, pKa, pKb, reaction rates, and equilibrium constants. In a particular embodiment, the partition coefficient is the octanol/water partition coefficient. Calculated non-spectral descriptors are one example of descriptors requiring structural knowledge beforehand that may be utilized along with segmented spectral data in establishing a predictive relationship for an endpoint.

Another aspect of the invention is a method of using spectral data as a set of structure descriptors for a compound that does not necessarily require knowledge of the compound's structure beforehand, by providing spectral data of a training set of compounds and segmenting the spectral data into bins.

Yet another aspect of the invention is a method for establishing a relationship between spectral data and a biological, chemical, or physical property, by providing spectral data for a training set of compounds, segmenting the spectral data into bins, and detecting patterns in the bins of the spectral data that are associated with the property. The method may also include detecting corresponding patterns in spectral data of test compounds to select the test compounds having the property. Furthermore, the test compounds may be

mixtures of compounds. The spectral data of the training set can be auto-scaled and weighted (for example by Fisher-weighting) to help identify data that are most strongly associated with the biological activity. Knowledge of the structural features that lead to the spectral data is not needed beforehand.

5 Yet another aspect of the invention is a method of determining the structural features of a plurality of compounds that contribute to determining a particular endpoint property exhibited by the compounds, by providing segmented spectral data for the plurality of compounds; providing endpoint data for the plurality of compounds; establishing a spectral data-activity relationship (SDAR)  
10 by identifying the segmented spectral features that bias toward increased endpoint values and the segmented spectral features that bias toward decreased endpoint values; and identifying the structural feature leading to the segmented spectral features that bias toward increased or decreased endpoint values for the plurality of compounds. As with the earlier described embodiments of the invention, the  
15 segmented spectral data may be a composite of several types of spectral data.

Any of the methods of the present invention can be performed without reference to a chemical structure of the test compound. Hence the spectral features of the training set and the test compound may be compared, without  
20 determining the chemical structure of the compounds of either the training set or the test compound or compounds. Any of the foregoing methods can also be incorporated into a computer readable medium, having stored thereon instructions for performing the steps of these methods.

The foregoing and other objects, features, and advantages of the invention will become more apparent from the following drawings and the detailed  
25 description of several embodiments.

### BRIEF DESCRIPTION OF THE DRAWINGS

Figs 1(a), 1(b) and 1(c) are the  $^{13}\text{C}$  NMR, EI MS, and IR spectra, respectively, for bisphenol A (4,4'-isopropylidenediphenol, structure shown in Fig.  
30 1(a)).

Figs. 2(a), 2(b) and 2(c) show tables of data that correspond respectively to each of the spectra shown in Figs. 1(a), 1(b) and 1(c).

Fig. 3 shows a hypothetical set of structure descriptors derived from the spectral data summarized in the tables of Figs. 2(a), 2(b) and 2(c).

5 Fig. 4 shows a flowchart of a particular embodiment of the spectral data-activity relationship (SDAR) method.

Fig. 5 shows the discriminant function using  $^{13}\text{C}$  Nuclear Magnetic Resonance (NMR) spectral data for 30 compounds in the Spectral Data-Activity Relationship (SDAR) model. The X-axis is the first canonical variate  
10 (discriminant function) and the Y-axis is the component frequency. The numbers in each box correspond to the numbers in Table 1 that serve to identify the compounds used in the SDAR. White boxes correspond to strong estrogen receptor binding compounds, and gray boxes correspond to medium estrogen receptor binding compounds.

15 Fig. 6 presents the first canonical variate factor weights using  $^{13}\text{C}$  NMR spectral data for 30 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. The bins are numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 7 shows the discriminant function using composite  $^{13}\text{C}$  NMR spectral  
20 data and Electron Impact (EI) mass spectral data for 30 compounds in the SDAR model. The X-axis is the first canonical variate and the Y-axis is the component frequency. The numbers in each box correspond to the numbers in Table 1 that serve to identify the compounds used in the SDAR. White boxes correspond to strong estrogen receptor binding compounds, and gray boxes correspond to  
25 medium estrogen receptor binding compounds.

Figs. 8(a) - 8(b) present the first canonical variate factor weights using composite  $^{13}\text{C}$  NMR spectral data and EI mass spectral data for 30 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. EI mass spectral data are in the bins numbered from m/z  
30 0 to 550.  $^{13}\text{C}$  NMR spectral data occupies the bins numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 9 shows the discriminant function using  $^{13}\text{C}$  Nuclear Magnetic Resonance (NMR) spectral data for 119 compounds in the Spectral Data-Activity Relationship (SDAR) model. The X-axis is the first canonical variate (discriminant function) and the Y-axis is the second canonical variate. The symbol S represents a strong estrogen receptor binder, M represents a medium estrogen receptor binder, and W represents a weak estrogen receptor binder.

Fig. 10 presents the first canonical variate factor weights using  $^{13}\text{C}$  NMR spectral data for 119 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. The bins are numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 11 shows the discriminant function using composite  $^{13}\text{C}$  NMR spectral data and Electron Impact (EI) mass spectral data for 119 compounds in the SDAR model. The X-axis is the first canonical variate and the Y-axis is the second canonical variate. The symbol S represents a strong estrogen receptor binder, M represents a medium estrogen receptor binder, and W represents a weak estrogen receptor binder.

Figs. 12(a) - 12(b) present the first canonical variate factor weights using composite  $^{13}\text{C}$  NMR spectral data and EI mass spectral data for 119 compounds in the SDAR model. The X-axis is the bin number and the Y-axis is the factor weight relative intensity. EI mass spectral data are in the bins numbered from m/z 0 to 550.  $^{13}\text{C}$  NMR spectral data occupies the bins numbered from 550 (corresponding to 0 ppm) to 770 (corresponding to 220 ppm).

Fig. 13 is a diagram of a distributed computing environment in which the present invention can be implemented.

Fig. 14 is a block diagram of a computer system that can be used to implement the present invention.

## DETAILED DESCRIPTION OF SEVERAL DISCLOSED EMBODIMENTS

The illustrative embodiments that follow are best understood by first understanding terms in the art that are relevant. These terms are defined below.

5

### Definitions and Abbreviations

**LOO** – leave-one-out cross validation

**RBA** – relative binding affinity

**NMR** – nuclear magnetic resonance

**IR**- infrared (spectroscopy)

10

**MS**-mass spectrometry

**EI MS** –electron impact mass spectrometry

**UV-Vis** –ultraviolet-visible (spectroscopy)

**CODESSA** – comprehensive descriptors for structural and statistical analysis

15

**SDAR** –spectral data-activity relationship

**Endpoint** – a particular biological, chemical, or physical property or a set of such properties for a compound that are either qualitatively or quantitatively measurable.

20

**Structure Descriptors** – any direct or indirect measure of the structure of a compound that may be obtained by theoretical or experimental means.

**Training Set** – endpoint data and structure descriptors (structural data) for a group of compounds used to establish a correlation between the endpoint property and the structures of the compounds.

25

**Validation Set** - endpoint data and structure descriptors (structural data) for a group of compounds used to test the reliability of the correlation between the endpoint property and the structures of the compounds.

30

**Segmented Spectral Data** – spectral data that is divided into discrete sub-spectral units (bins), each of which spans a particular spectral range. The spectral range spanned by a particular bin corresponds to a range of frequencies or a range of wavelengths for spectroscopic data and may be equal to the digital resolution of the spectral data or greater. For mass spectrometric techniques, the spectral range



within each bin corresponds to a particular mass or range of masses and may be equal to the digital resolution of the spectral data or greater. The bins need not all be of equal width. The spectral data that is divided into bins may either encompass all the spectral data of a particular type that are available or cover only a portion of the spectral data of a particular type that are available. Each bin contains information derived from the spectral signals (or lack thereof) that appear within the spectral range defined by a particular bin. The structural aspect(s) of the compounds that give rise to the information falling within any particular bin need not be known.

10       **Spectral Data-Activity Relationship (SDAR)** – a correlation between endpoint data and spectral data for a group of compounds, useful for among other things for predicting the endpoint data for compounds from their spectral data.

**Nuclear Magnetic Resonance (NMR)** – a phenomenon exhibited by a large number of atomic nuclei in which nuclei in a magnetic field absorb energy from a radio-frequency field at certain characteristic frequencies. Particular examples of nuclei that exhibit this phenomenon include  $^{13}\text{C}$ ,  $^1\text{H}$  and  $^{31}\text{P}$ .

**Mass Spectrometry (MS)** – a method of chemical analysis in which the substance to be analyzed is placed in a vacuum and reduced to low pressure. The resulting vapor is exposed, for example, to a beam of electrons which causes ionization to occur, either of the molecules or their fragments. The ions thus produced are accelerated and then passed through a mass analyzer that separates the ions according to their mass.

**Electron Impact Mass Spectrometry (EI MS)** – a mass spectrometric technique in which the ionization of molecules and their fragments is accomplished by a beam of electrons that impacts the molecules and their fragments. Typically, as the energy of the electron beam is increased, the number of fragments produced from a molecule increases.

**Infrared Spectroscopy (IR)** – an analytical technique which measures a range of wavelengths (or frequencies) in the infrared region or near-infrared region of the electromagnetic spectrum that are absorbed by a specimen, which characterize its molecular constitution. Infrared absorption bands identify

molecular structure components, such as aromatic, olefin, aliphatic, aldehyde, ketone, carboxylic acid, alcohol, amine, and amide groups. The frequency at which absorption occurs also reflects the frequency at which the bonds in these components stretch and and/or bend.

5           **Ultraviolet-Visible Spectroscopy (UV-Vis)** – an analytical technique which measures a range of wavelengths (or frequencies) in the ultraviolet and visible regions of the electromagnetic spectrum that are absorbed by a specimen, which characterize the electronic energy levels of its molecular constituents. UV-Vis absorption bands may be characteristic of certain molecular components, such as  
10 aromatic groups or carboxyl (CO) groups.

**Fluorescence Spectroscopy** – an analytical technique which measures a range of wavelengths (or frequencies) of light a molecule emits in passing from a higher to lower energy electronic state during a given time period (such as the first millisecond) after absorbing a photon of light. Fluorescence wavelengths and  
15 emission intensity reflect the redistribution of energy in the molecule after light absorption. Fluorescence excitation spectroscopy reflects the efficiency with which a molecule converts absorbed energy into fluorescent emission as a function of the wavelength of the absorbed photons.

**Phosphorescence Spectroscopy** - an analytical technique which measures a  
20 range of wavelengths (or frequencies) of light a molecule emits in passing from a higher to lower energy electronic state on a time scale beyond the first millisecond after absorbing a photon of light. Phosphorescence wavelengths and emission intensity also reflect the redistribution of energy in the molecule after light absorption. Phosphorescence excitation spectra reflect the efficiency with which a  
25 molecule converts absorbed energy into phosphorescent emission as a function of the wavelength of the absorbed photons.

**Principal Component Analysis (PCA)** – this pattern recognition technique and the related technique of partial least squares analysis reduce the structure descriptors used in SDAR to a smaller number of linear combinations of the  
30 original set of descriptors that are called principal components (PCs). The optimum number of principal components will yield the smallest standard error of

prediction while capturing as much of the variance shown by the structural data as possible. (See generally, Kramer, R., *Chemometric Techniques for Quantitative Analysis*, Marcel Dekker, Inc., 1998)

**Comprehensive Descriptors for Structural and Statistical Analysis**

- 5 (CODESSA) – a set of various structural descriptors typically utilized in three dimensional quantitative structure-activity studies that includes constitutional descriptors, topological descriptors, geometrical descriptors, electrostatic descriptors, and quantum mechanical descriptors, all of which require knowledge of structure beforehand. (See , Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995  
10 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference)

- Auto-scaling** – a method whereby the quantitative spectral information contained within each particular bin is compared for all compounds in the training set to yield an average value and a standard deviation. Then, for each bin  
15 comprising the structure descriptors of a given compound, the quantitative spectral information therein is expressed as a number of standard deviations above or below the average for each bin. Autoscaling equalizes the importance of inherently weak spectral signals falling within certain bins with the importance of inherently strong spectral signals falling within certain other bins in describing a  
20 set of spectrally derived structure descriptors. It may also equalize the importance of different types of spectral data in a composite of spectral data.

- Fisher-weighting** – a method whereby the quantitative spectral information in bins that are important for classifying the training set compounds into different endpoint groups, such as strong and medium binders to the estrogen receptor, are  
25 enhanced. For each bin, the variance of the quantitative spectral information between the endpoint groups is divided by the variance of the quantitative spectral information within the endpoint groups. The resulting dividend becomes a weighting factor that has a magnitude larger than one when a particular bin has an important role in distinguishing the endpoint groups. Each bin is multiplied by its  
30 weighting factor to yield structure descriptors that are more sensitive to subtle but significant spectral variations.

**Leave-one-out (LOO) Cross-Validation** - a method whereby each compound in the training set is systematically excluded from the data set, after which its endpoint value is predicted by the spectral data-activity relationship derived from the remaining compounds (See, Cramer et al., *Quant. Struct-Act. Relat.* 7: 18-25, 1998, incorporated herein by reference). Cross-validation is useful for judging the reliability of a spectral data-activity relationship, especially where a validation set of compounds is not available.

### EXAMPLES

The following examples further illustrate the QSAR methods of the present invention. The methods utilize spectral data as structure descriptors and correlate the spectral data with specific biological, chemical, or physical endpoints without the need to assign spectral features to their corresponding structural elements. In recognition of the advantages presented by using spectral data as structure descriptors, a correlation provided by the methods described herein is termed a Spectral Data Activity Relationship (SDAR).

In various embodiments, experimental and/or calculated molecular spectral data may be used as structure descriptors. If experimental spectral data is used in a particular embodiment, there is no need to actually know the molecular structures beforehand. Calculated molecular spectral data may be utilized as a surrogate for experimental data in some embodiments. In other embodiments, calculated molecular spectral data may be useful, in combination with an SDAR derived from experimental spectral data, for screening compounds that are created as part of a computer generated combinatorial library. In yet other embodiments, calculated molecular spectral data may be generated for compounds for which no experimental spectral data is available and used to establish an SDAR or to screen those compounds for a particular property.

In some embodiments, spectral data is used as a set of descriptors, for example descriptors of molecular structure. The pattern of the spectrum is determined, for example by segmenting the spectral data into portions covering particular spectral regions (e.g., ranges of frequency, wavelength, chemical shift,

mass to charge ratio, and the like). The number and/or the intensity of the spectral signals within each segmented region may serve as the structure descriptors. The spectral data may be of one type or be a composite of several types, including NMR, MS, IR, UV-Vis, Fluorescence, and Phosphorescence. Spectral data of a particular type may be utilized in its entirety or in part. While spectral data are often used to elucidate the structure of the compound that yields them, the information contained in the spectra may be used in some embodiments without the need to interpret the spectra. Furthermore, the spectral data may be used in certain embodiments without the need to know the structures of the compounds beforehand. Segmented spectral data is particularly amenable to encryption for secure analysis.

In some embodiments,  $^{13}\text{C}$  NMR, EI MS, and IR spectra may be combined to yield a set of structure descriptors. Examples of these types of spectra for the compound bisphenol A are shown in Figs. 1(a), 1(b), and 1(c) respectively. Each type of spectrum contributes information regarding the structure and electronic properties of a molecule. For example, in the  $^{13}\text{C}$  NMR spectrum shown in Fig. 1(a), the peaks reflect the number of different types of carbon atoms and the electronic environment of each type of carbon atom in bisphenol A. Peaks exhibiting the largest  $\delta$  (chemical shift) values correspond to carbon atoms of a type that are the least shielded from the applied magnetic field by the electrons of the molecule. The EI MS spectrum shown in Fig. 1(b) reveals the molecular mass ( $m/z = 228$ , 228 amu) of bisphenol A and the mass of the fragments into which the molecule breaks up when the electrons impacting the molecule deliver enough energy to break bonds. For example, the most prominent peak in the mass spectrum shown in Fig. 1 (b) appears at a  $m/z$  of 213. The peak at  $m/z=213$  corresponds to the fragment formed by the loss of a methyl group ( $\text{CH}_3$ ,  $m/z = 15$ ) from bisphenol A (i.e.  $228-213 = 15$ ). The mass spectral peaks thus reflect the fragments of structure that comprise the molecule.

The IR spectrum shown in Fig. 1(c) reveals the various types of bonds that are present in bisphenol A. For example the peak that appears at  $3368\text{ cm}^{-1}$  in the IR spectrum corresponds to the two oxygen-hydrogen bonds found in bisphenol A.

While the  $^{13}\text{C}$  NMR, EI MS and IR spectral features have been discussed with reference to the structure of bisphenol A to illustrate the types of structural information contained in these spectra, it should be noted that in some embodiments the structure of the molecule responsible for the features need not be known and no attempt need be made to assign particular spectral features to corresponding structural features.

The spectral data from Figs. 1(a), 1(b) and 1(c) are compiled in tabular form in Figs. 2(a), 2(b) and 2(c), respectively, as tables of spectral frequencies and relative intensities. Such compilations are typical of the output of modern digital spectrometers. The data as presented in these tables may be used to construct a set of molecular structure descriptors for bisphenol A and similar sets of data may be used to generate structure descriptors for any number of compounds.

To construct the set of structure descriptors shown in Fig. 3, each type of spectrum is segmented into spectral "bins" and the information content of each bin becomes a separate structure descriptor. Fig. 3 shows a set of raw structure descriptors for bisphenol A derived, for illustrative purposes only, from the spectral data as presented in the tables of Figs. 2(a), 2(b) and 2(c). It is possible to include more information from the spectra than considered here, such as the intensity of IR absorption at frequencies that do not correspond to a distinct peak. Each bin in this case corresponds to a particular integer  $m/z$  ratio for the EI MS data, a range of chemical shift frequencies for the  $^{13}\text{C}$  NMR data, and a range of wavenumber frequencies for the IR data. Moreover, each bin contains the relative intensity of the spectral feature(s) falling within a range covered by a particular bin. In this example, bins 1 to 49 were not used (because such low mass ions are common and may also result from air contamination), bins 50 to 549 correspond to  $m/z$  ratios from 50 to 549, bins 550 to 770 correspond to 1 ppm segments of the  $^{13}\text{C}$  NMR spectrum over the range of 0 to 221 ppm, and bins 771 to 1120 correspond to  $10\text{ cm}^{-1}$  segments of the IR spectrum over the range from  $4000\text{ cm}^{-1}$  to  $500\text{ cm}^{-1}$ . Those spectral bins that do not contain spectral data are given a value of zero and are not shown in Fig. 3. Other regions of the spectra can be selected to correspond to the bins in other embodiments of the invention, which is not

limited to the specific examples recited herein. In addition, the spectra can be divided into different segments, instead of the integer segments that are merely given as examples.

Tracking specific spectral data from the various spectra to the set of structure descriptors in Fig. 3 helps to illustrate how, in some embodiments, the spectral data is utilized to construct the structure descriptors. For example, the intensity of the peak at  $m/z=213$  in the mass spectrum appears in bin number 213. The intensity of the peak appearing at 154.2 ppm in the  $^{13}\text{C}$  NMR spectrum is in bin number 704 because the peak falls within the range of 154 ppm to 155 ppm which corresponds to bin number 704. The peak appearing at  $3368\text{ cm}^{-1}$  in the IR spectrum falls within the frequency range  $3370\text{ cm}^{-1}$  to  $3360\text{ cm}^{-1}$  and thus the intensity of the peak appears in bin number 834 since this bin corresponds to this particular frequency range. These three entries are highlighted and shown in bold in Fig. 3.

The segmented spectral data structure descriptor set for bisphenol A shown in Fig. 3 may be altered in some embodiments to normalize the data from each type of spectrum to yield structure descriptors of similar magnitude from each type of spectral data. For example, in order to normalize the IR data and  $^{13}\text{C}$  NMR data to the greatest value in the MS data (100), they might be expressed as the number of peaks falling within a particular bin's range, multiplied by 100. The 21 appearing in bin 834 would thus be replaced with the value 100 as would the 411 appearing in bin 704, because in both cases only one peak appears in the spectral ranges corresponding to these bins.

In other embodiments, UV-Vis, fluorescence, and phosphorescence spectral data is segmented into bins alone or in combination with each other and with NMR, MS, and IR data.

In addition to not requiring structures beforehand, experimental spectral data as used in some embodiments offers several advantages over calculated spectral data. Most experimental spectral data reflects the quantum mechanical properties of molecules as determined by both structure and solvation. However, mass spectra do not reflect solvation. Experimental spectral data also avoids the

difficulty and approximations inherent in deriving structure descriptors from quantum mechanical or electrostatic potential calculations, which are especially inaccurate when the calculations attempt to include solvation effects. Experimental spectral data may be reflective of the solution conformations that are responsible for a particular endpoint property as well as the role molecular flexibility plays in determining the endpoint property. Moreover, experimental spectral data reflect specific structural features and are therefore preferable to bulk structure descriptors such as partition coefficients that lack structural specificity. Additionally, experimental spectral data is already available for many compounds in various databases.

In some embodiments, segmented spectral data is further pre-treated before being combined with endpoint data and subjected to statistical pattern recognition or artificial intelligence based pattern recognition to extract an SDAR. Fig. 4 summarizes, in flowchart form, a particular embodiment of the steps that may be taken to establish an SDAR from spectral data. These steps and others may be performed using a computer system.

In Fig. 4, spectral data is obtained at 20, and then the spectral data is segmented into bins at 22, which in some embodiments is carried out in a manner similar to that described above with reference to Figures 1, 2, and 3. The process of segregation is repeated for the spectral data of each of the compounds in the training set. The spectral data for each of the compounds of the training set can be segmented into a matrix of bins that represent the training set. These segmented spectral data sets 24 comprise the structure descriptors for the training set compounds. Patterns in the matrix of bins can then be determined, for example by pattern recognition software.

In some embodiments, the segmented spectral data 24 is pretreated by auto-scaling 26 prior to pattern recognition, to produce auto-scaled, segmented spectral data 28. Auto-scaling is a method whereby the quantitative spectral information contained within each particular bin is averaged over all the compounds in the training set to yield an average value and a standard deviation for each bin. Then, for each bin comprising the structure descriptors of a given compound, the



quantitative spectral information therein is expressed as the number of standard deviations above or below the average value for that bin. This reduces the variation in numerical magnitude between the bins in the segmented spectral data. Auto-scaling thus helps equalize the importance of inherently weak spectral signals that fall within the spectral range of certain bins with the importance of inherently strong spectral signals that fall within the spectral range of other bins.

In some embodiments, the segmented spectral data is further pre-treated by Fisher-weighting 30 the auto-scaled data 28 to improve the ability of pattern recognition algorithms to discern which bins (and the spectral information corresponding to the spectral range covered by the bin) are most important for classifying compounds into two or more endpoint classes (such as strong and weak estrogen receptor binders). An endpoint class is assigned to each compound based upon its endpoint value. For example, compounds could be classified into two endpoint classes, those compounds that have an endpoint value above a certain number and those that have an endpoint value equal to or below a certain number. For each bin of the segmented spectral data 24 the variance in value between the endpoint classes for the bin is divided by the variance in value within the endpoint classes for the bin to yield a weighting factor. The weighting factor has a magnitude larger than one when a particular bin has an important role in distinguishing the endpoint classes.

Fisher-weighting 30 quantifies the tendency for certain bins (and the spectral range corresponding to the bin) to be more helpful than others in deciding the endpoint class of a compound. For example, consider a particular bin in which the value in the bin is always large for the compounds belonging to one endpoint class (such as strong estrogen receptor binders), and always small for the compounds of a second endpoint class (such as weak estrogen receptor binders). Knowing that the spectral data for a compound yields a large value in that particular bin helps to classify the compound into the first endpoint class. If on the other hand, the value found in the bin is always large for compounds of both classes, it reveals nothing about the endpoint class of the compound to know that it has a large value in the bin. In the first case, the variance between the classes

would be larger than the variance within each class and the weighting factor will be greater than one. Conversely, in the second case, the variance between the classes will be about equal to the variance within the class, making the weighting factor equal to about one. Each bin of the segmented spectral data is then

5 multiplied by its weighting factor to yield a set of structure descriptors 32 that emphasize the bins most important for deciding the class of a compound. Such data is more easily treated by pattern-recognition analysis.

Pattern recognition 34 is used to establish an SDAR 36, by correlating the segmented (and optionally pretreated) spectral data with the endpoint data for the

10 compounds in the training set. In particular embodiments, the compounds are classified into two or more endpoint classes (e.g. strong versus weak estrogen receptor binders) according to their relative endpoint values. The pattern recognition then determines any segmented spectral features (i.e., the bins) that are characteristic of the compounds falling into each of the classes. If Fisher-

15 weighting is performed prior to pattern recognition, the endpoint classification scheme used for Fisher-weighting may be retained during pattern recognition. A simplified example is that a bin with a strong signal in all the compounds of the test set that have strong estrogen receptor binding would be a bin that would be considered as predictive of strong estrogen receptor binding if a test compound

20 also exhibited a strong signal in that bin. Multiple such signals would, however, usually be taken into account by the pattern recognition software when determining spectral patterns that are associated with the endpoint.

Pattern recognition to establish an SDAR 36 may be accomplished by statistical methods or artificial intelligence methods. Underlying these methods is

25 the idea that if a particular bin corresponds to a spectral signal exhibited by only the compounds of a particular class (e.g. strong estrogen receptor binders), the bin will bias strongly toward that class. The extent to which a particular bin biases toward a particular class depends upon whether or not compounds from all classes (e.g. strong and weak estrogen receptor binders) exhibit spectral signals

30 corresponding to the bin. For example, if half of the compounds in one class show a signal that falls within a particular bin and slightly more than half of the

compounds in another class show a signal in the same bin, the bin may only bias slightly toward the latter class. Statistical and artificial intelligence methods attempt to quantify these biases and provide a basis for classifying compounds to an endpoint class according to their segmented spectral data.

5           Once quantified, the relationship between the segmented spectral data and the endpoint may be visualized in various ways, depending upon the particular software package utilized for pattern recognition. One way to visualize the extent to which individual bins of segmented spectral data bias toward endpoint classes is with a canonical variate factor plot. An example of a canonical variate factor plot  
10 is shown in Fig. 6, which is discussed further with respect to Example 1. The SDAR model used to generate Fig. 6 was based upon two endpoint classes. The peaks pointing in one direction bias toward one class (strong estrogen receptor binding) and the peaks pointing in the other direction bias toward the other class (moderate estrogen receptor binding). The length of the peaks in Fig. 6  
15 correspond to how strongly the peaks bias toward a particular endpoint class. For example, the peak appearing in bin 585 of Fig. 6 strongly biases toward the class of strong binders while the peak appearing in bin 579 of Fig. 6 biases strongly toward the other class of moderate binders.

          An SDAR, once established by pattern recognition, may be used in some  
20 embodiments to quantitatively predict the endpoint value and endpoint class of any compound according to its segmented spectral data. In a particular embodiment, segmented spectral data of the type utilized to establish the SDAR is considered along with the canonical variate function to yield a prediction of endpoint class. For example, in reference to Fig. 6, a compound may be qualitatively predicted to  
25 belong to the class indicated by the upward pointing peaks if it exhibits a large number of spectral features falling into the upward biasing bins. The SDAR quantifies the prediction by considering not only the number of spectral features that fall into upward or downward biasing bins, but the strength with which each bin biases toward a particular endpoint class.

30           One example of a way to visualize the ability of an SDAR to correctly classify compounds into their endpoint class is with a discriminant function plot.

An example of a discriminant function plot is given in Figure 5, which is discussed in more detail with respect to Example 1. This discriminant function plot is also based upon the same two class SDAR used to generate the canonical variate function of Fig. 6. The grey and white boxes correspond to the two endpoint classes (e.g. strong and moderate estrogen receptor binding) into which the training set data was divided. The aggregation of grey boxes and the aggregation of the white boxes, as well as the separation of the two aggregates, illustrate the ability of the SDAR to discriminate between the two endpoint classes. If the SDAR had not found a correlation between the segmented spectral data and the endpoint classes, the discriminant function would have exhibited a greater mixing of the grey and white boxes.

Unlike methods using spectral data that is referenced to a parent compound's spectral data, the methods of the present invention can in certain embodiments reveal the importance of particular structural features in determining the endpoint value without requiring knowledge of the identity of those structural features beforehand. For example, if a particular bin is identified by the canonical variates as showing a bias toward strong anti-tumor activity, subsequent elucidation of the structure responsible for the signal occupying that particular bin is possible. Furthermore, spectral data as used in some embodiments is well suited for examining structure activity relationships for compounds of diverse structures where several different groupings of atoms may elicit similar quantum mechanical features that are important to a particular endpoint property.

Another benefit of using spectral data to establish an SDAR according to the methods of the present invention is the flexibility it offers. Once spectral data are gathered for a set of compounds, an SDAR may be generated with reference to a multitude of biological, chemical, or physical endpoints for which data is  
5 available. For instance, the same set of spectrally derived structure descriptors for a set of compounds may be utilized with toxicity data and antibacterial activity data for these compounds to establish two separate SDARs for the compounds, one for toxicity and the other for antibacterial activity.

Because the experimentally based SDAR method does not require input of a  
10 compound's structure, the method yields accurate results without computer modeling of a compound's steric or electrostatic properties. The method also removes the need to break the compound into secondary structural motifs or assume a particular alignment for how a set of molecules will bind to another molecule.

15 Methods according to the invention additionally provide rapid and inexpensive approaches to screen compounds for selected activities or properties. Screening using these methods does not require that time and effort be expended in trying to elucidate the structure of compounds beforehand, like screening using 3D-QSAR. Screening according to the methods of the present invention, by  
20 measuring a compound's spectral data and inputting it into an SDAR, is much more rapid than screening with bioassays. Furthermore, once obtained, the same spectral data for a compound may be utilized in multiple SDARs to predict various properties of the molecule. Additionally, the methods may also be used to screen mixtures or fractions of compounds for spectral features associated with a  
25 particular property, thereby obviating the need to spend time and money isolating compounds that show no promise of having the particular property. The methods may therefore be extremely useful, for example, in combination with experimentally generated combinatorial libraries of compounds.

When implemented in a real-time web-based computer environment, the  
30 methods of the present invention may provide investigators with a means for rapidly and securely estimating the molecular activities of proprietary molecular

structures. Spectral data as utilized in the present invention may be generated for any number of compounds and submitted by an investigator, either directly or as a segmented set of descriptors, to a central location containing the SDAR software. The spectral data or segmented spectral data may be encrypted by the user and

5 securely transmitted to the central location. Because of the speed at which pattern recognition software can make predictions based upon an SDAR, a submission of spectral data and retrieval of a prediction based on an SDAR can take place in real time over the worldwide web without the spectral data being stored at the central location. Alternatively, the SDAR and the pattern recognition software may be

10 implemented on a single computer in an investigator's laboratory. In either case, there is no need to reveal the structure of a proprietary compound to obtain a prediction. The ability to provide spectral data in a pre-segmented and encrypted form that is difficult to use for structure elucidation purposes (i.e., industrial espionage) may also encourage investigators to submit their spectral and endpoint

15 data permanently to a central location for use as part of larger training sets that may lead to more reliable SDARs.

It will be understood, however, that many variations and modifications of the methods can be made while remaining within the scope and spirit of the present invention. For example, segmentation of spectral data into bins need not occur,

20 but is only one example of how spectral patterns may be observed and compared. Other approaches for comparing spectral patterns could also be used, for example artificial neural networks that may not require gross segmentation of the spectral data per se (image analysis).

25 **Example 1- Use of  $^{13}\text{C}$  NMR and EI Mass Spectrometric Data to Produce a Predictive Model of Estrogen Receptor Binding Activity.**

The Food Quality Protection Act, passed in 1996, mandates that the Environmental Protection Agency (EPA) develop screening and testing procedures for endocrine disrupting chemicals (EDCs). An EDC is defined as "an exogenous

30 agent that interferes with the production, release, transport, metabolism, binding, action, or elimination of natural hormones in the body responsible for the

maintenance of homeostasis and the regulation of developmental processes.”

Estrogenic compounds represent a significant subset of the EDCs to be tested.

Many of these compounds can be screened by determining how strongly the compounds bind to estrogen receptors. Thus there is a need to develop

5 inexpensive and rapid methods to screen thousands of potential compounds.

SDAR according to the invention meets this need.

The  $^{13}\text{C}$  nuclear magnetic resonance (NMR) spectrum of a compound contains frequencies that correspond directly to the quantum mechanical properties of the molecule and depend largely on the electrostatic features and geometry, including the stereochemical configuration, of the molecule. Furthermore, a  
10 solution spectrum, like NMR, inherently reflects the effects of solvation on the quantum mechanical properties. Electron-impact mass spectral (EI MS) data provide a mass-size description of molecular substructures (and possibly the whole molecule) as well as information about the strength of bonds between the atoms of  
15 the molecules. The  $^{13}\text{C}$  NMR and EI mass spectral data thus represent sets of quantum mechanical structure descriptors that reflect the electrostatic and steric properties of a molecule. Such experimental data are readily attainable and often already available.

Two Spectral Data-Activity Relationship (SDAR) models were established, based on  $^{13}\text{C}$  NMR and EI MS spectra, for 30 estrogenic chemicals whose relative  
20 binding affinities (RBA) for the alpha ( $\text{ER}\alpha$ ) and beta ( $\text{ER}\beta$ ) estrogen receptors are available. The SDAR models classified the 30 compounds into two endpoint classes, those with strong binding affinities for the estrogen receptors and those with medium binding affinities for the estrogen receptors. The SDAR based on  
25  $^{13}\text{C}$  NMR data alone yielded a leave-one-out cross-validation of 90% and the SDAR based on a composite of  $^{13}\text{C}$  NMR and EI MS data yielded a cross-validation of 83%. Two compounds that were classified incorrectly in both SDAR models were in a transition zone between these classifications. Both SDAR models correctly classified 25 compounds. The models provided rapid, reliable,  
30 and simple ways to predict whether a compound binds to the estrogen receptor.

### Procedure

The estrogenic relative binding affinities (RBAs) of the 30 compounds were obtained from previous publications (Kuiper et al., *Endocrinology* **138**:863-870, 1997; and Tong et al., *Endocrinology*, **138**:4022-4025, 1997). Most of the  $^{13}\text{C}$  NMR spectrometric and EI mass spectrometric data were obtained from the Integrated Spectral Data Base System for Organic Compounds web site [www.aist.go.jp/RIODB/SDBS](http://www.aist.go.jp/RIODB/SDBS), the *Aldrich Library of  $^{13}\text{C}$  and  $^1\text{H}$  FT NMR Spectra* (Pouchert and Behnke, Eds., Aldrich Chemical Company, Volumes 1-3, 1993), *Spectral Data of Steroids* (Frenkel and Marsh, eds., Thermodynamics Research Center: College Station, 1994), and the NIST MS database software version 1.6. Experimental  $^{13}\text{C}$  NMR and EI MS data for five compounds were obtained using standard methods.

Specifically, the  $^{13}\text{C}$  nuclear magnetic resonance (NMR) spectral analyses of 4-hydroxy-estradiol, ICI 164,384, moxestrol, and norethynodrel were performed at 75.46 MHz on a Varian Gemini 300 MHz NMR (Varian Associates, Inc., Palo Alto, CA) spectrometer operating at 301 K. The subject compounds were dissolved in  $\text{CDCl}_3$  or DMSO (dimethyl sulfoxide). The chemical shifts were defined by denoting the  $\text{CDCl}_3$  peak at 77.0 ppm and the DMSO peak at 39.5 ppm. The spectral width was 21,008 Hz with a 2.6-second delay time between acquisitions. The acquisition time was 0.495 seconds and the number of points acquired was 20,800.

The samples were also analyzed by direct exposure probe (DEP) mass spectrometry (MS). The mass spectrometers were operated in the electron-impact (EI) mode, with 70 V electron energy. The ion source temperature was set at 150°C. Samples, in solution, were applied to the rhenium wire of the DEP and the solvent was allowed to evaporate before the analysis was begun. MS data were collected until the current of the DEP exceeded 500 mA. Norethynodrel, ICI 164,384, 4-hydroxy-tomoxifen, and 4-hydroxy-estradiol were analyzed on a Finnigan TSQ 700 mass spectrometer and moxestrol was analyzed on a Finnigan 4500 mass spectrometer (Finnigan Corp., San Jose CA).



The  $^{13}\text{C}$  NMR and EI-mass spectral data points were used as comprehensive descriptors for structural and statistical analyses (CODESSA) as in Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference). Mass spectrometric data for mass-to-charge ( $m/z$ ) ratios from 50 to 549 were used directly as ordered pairs of  $m/z$  and relative intensity (maximum of 100). Unassigned 1D  $^{13}\text{C}$  NMR chemical shifts were segmented into bins over a 0 to 221 ppm range. The  $^{13}\text{C}$  NMR frequencies were shifted to bins 550 to 770, so bin 550 was the  $^{13}\text{C}$  NMR spectrum for frequencies inside 0 to 1 ppm and bin 770 was the  $^{13}\text{C}$  NMR spectrum for frequencies inside 220 to 221 ppm.

To save space, the  $^{13}\text{C}$  NMR spectra were saved as sets of ordered pairs each consisting of the respective chemical shift frequency in ppm and the respective area under the peak. The area under the peak of a specific chemical shift frequency was first normalized to an integer. A non-degenerate frequency was assigned an area of 1; a doubly degenerate frequency had an area of 2; and so forth. This was done to provide all the spectra with a similar signal-to-noise ratio and to eliminate line-width variations due to differences in NMR-instrument field strengths, shimming, temperature, pH, and solvents. The bin defined the number of significant and distinct chemical-shift peaks inside a ppm range. The optimal range in this example was found to be 1 ppm to each bin. To normalize the peak intensity of the  $^{13}\text{C}$  NMR spectroscopic data relative to the mass spectrometric data, the area of one  $^{13}\text{C}$  NMR peak was set to be 25. The number 25 was selected with the objective of scaling the maximum value for the  $^{13}\text{C}$  NMR data inside a bin to near 100, which is the maximum value for EI MS data. The number of bins used to input the  $^{13}\text{C}$  NMR spectra was studied by varying the width of the bins from 0.5 to 5.0 ppm. Again, the optimal ppm range for this study was found to be 1.0 ppm. Values were normalized to a maximum of 100 prior to pattern-recognition analysis.

The relative binding affinity (RBA) to the estrogen receptor was defined as the ratio of the molar concentration of 17- $\beta$ -estradiol to the competing compound required to decrease the receptor-bound 17- $\beta$ -estradiol by 50%, multiplied by 100.

Thus, 17- $\beta$ -estradiol has an RBA of 100 and a log (RBA) of 2.0. Strong binders to the estrogen receptor were classified as having a log (RBA) over 0.0, and medium binders to the estrogen receptor were classified as having a log (RBA) of less than or equal to 0.0. There were 17 strong binders and 13 medium binders in the training set. No weak estrogen-receptor-binding compounds (i.e., those with a log (RBA) of less than or equal to -3) were included. SDAR models that include more compounds, including weak-binding compounds, in the training set are better able to accurately predict strong estrogen receptor binding (see example 2).

The analysis of the SDAR model was performed by the leave-one-out (LOO) cross-validation procedure in which each compound was systematically excluded from the training set and its relative binding activity class predicted by the model (Cramer et al., *Quant. Struct.-Act. Relat.*, 7:18-25, 1988, incorporated by reference).

The pattern-recognition software used was RESolve Version 1.2 (Colorado School of Mines, Boulder, CO). The  $^{13}\text{C}$  NMR and EI MS spectroscopic data for all 30 compounds were input as text files into a computer programmed with the software. The spectroscopic data was then auto-scaled and Fisher-weighted prior to principal component analysis (PCA). The discriminant analysis was based on canonical variate vectors. Leave-one-out (LOO) cross-validation was used to maximize the size of the training set.

Autoscaling compared the quantitative response at each mass spectral m/z bin or NMR chemical shift bin to all the other bins in the comparison set. An average value (with standard deviation) was calculated for each bin. Then, for each spectrum, the quantitative response at each bin was expressed as the number of standard deviations above or below the respective average. This data-pretreatment step equalized the weight of consistent variance of signals with inherently small magnitudes (25 units for the NMR bin 558 representing a single methyl carbon) to those signals with large magnitudes (130,000 area counts at m/z bin 91, possibly arising from a tropylium ion fragment). Autoscaling automatically compensates for gross magnitude variations. It was particularly helpful in this example in which two completely different types of analytical

spectra were formed into a composite set of structure descriptors that are representative of molecular characteristics.

Fisher weighting was used to emphasize those spectral characteristics important in distinguishing defined endpoint groups. For each mass spectral m/z or NMR bin, the variance between groups was divided by the variance within groups. The resulting dividend became a weighting factor with a magnitude larger than one when the particular m/z or bin has an important role in distinguishing groups. Fisher weighting all of the raw spectra before pattern recognition increased the power of discriminant analysis to classify spectra correctly. In this example, Fisher weighting de-emphasized the relative importance of irrelevant spectral information (such as the NMR signals from carbon atoms not important to binding to the estrogen receptor).

## Results

Based only on  $^{13}\text{C}$  NMR spectroscopic data, the statistical pattern-recognition program with 8 principal components (PCs) had 81.1% of the total variance and a cross validation of 90%. The mathematical mode of significance was 91.9% at 1, with 28 degrees of freedom. The Wilks discriminant criterion was 99.99%. Table 1 lists the compound names, an identifying number for each compound that is used in Figs. 5 and 7, the log (RBA) alpha and log (RBA) beta values, the SDAR training input endpoint class for each compound ("class input")(either S for strong binding or M for medium binding), the SDAR prediction ("class predicted NMR") from the  $^{13}\text{C}$  NMR data alone and the SDAR prediction ("class predicted NMR and MS") from the  $^{13}\text{C}$  NMR and EI MS data. Fig. 5 shows the discriminant function for SDAR derived from the  $^{13}\text{C}$  NMR data alone which illustrates the ability of the SDAR to discriminate between the two endpoint classes in the training set. Compounds denoted with a white background exhibit strong RBAs, and compounds with a gray background exhibit medium RBAs. Fig. 5 shows that the SDAR yielded a large separation between the 15 strong (white rectangles on right side) and the 9 medium binders (gray rectangles on the left side); 4 strong and 2 medium RBAs are in a transition zone (middle)

between strong and medium RBAs. Table 1 also shows that 27 of the 30 compounds are correctly group-predicted using only,  $^{13}\text{C}$  NMR data.  $3\beta$ -androstenediol, 2-hydroxy-estradiol and hexestrol are incorrectly predicted to have a medium RBA using 8 principal components. Only  $3\beta$ -androstenediol and

5 hexestrol are incorrectly predicted to have a medium RBA using the canonical variate function. Strong binding  $3\beta$ -androstenediol is most likely incorrectly predicted because the compound is similar to the medium RBA  $3\alpha$ -androstenediol. Repeating the analysis using a larger training set (Example 2) eliminated even this confusion.

TABLE 1

COMPOUND	Number	Log(RBA) alpha	Log(RBA) beta	Class Input	Class Predicted NMR	Class Predicted NMR + MS
Coumestrol	1	1.97	2.27	S	S	S
3 $\alpha$ -Androstenediol	2	-1.15	-0.52	M	M	S
3 $\beta$ -Androstenediol	3	0.48	0.85	S	M	M
4-OH-Estradiol	4	1.11	0.85	S	S	S
17 $\alpha$ -Estradiol	5	1.76	1.04	S	S	S
17 $\beta$ -Estradiol	6	2.00	2.00	S	S	S
Bisphenol A	7	-1.30	-0.48	M	M	M
$\beta$ -Zearanol	8	-1.20	-1.15	M	M	M
Clomifene	9	1.40	1.08	S	S	S
2-OH-Estradiol	10	0.85	1.04	S	M	S
Dehydroepiandrosterone	11	-1.40	-1.15	M	M	M
Diethystibesterol	12	2.67	2.47	S	S	S
Dienestrol	13	2.35	2.61	S	S	S
Dihydrotestosterone	14	-1.30	-0.77	M	M	M
Estriol	15	1.15	1.32	S	S	S
Estrone	16	1.78	1.57	S	S	S
Genistein	17	0.70	1.56	S	S	S
Hexestrol	18	2.48	2.37	S	M	M
ICI 164,384	19	1.93	2.22	S	S	S
Methoxychlor	20	-2.00	-0.89	M	M	M
Moxestrol	21	1.63	0.70	S	S	S
Nafoxidine	22	1.64	1.20	S	S	S
Nandrolone	23	-2.00	-0.64	M	M	M
Norethindrone	24	-1.15	-2.00	M	M	M
Norethynodrel	25	-0.16	-0.66	M	M	M
Progesterone	26	-3.50	-3.50	M	M	M
Tamoxifen	27	0.85	0.78	S	S	S
Testosterone	28	-2.10	-2.10	M	M	M
5 $\alpha$ -Androstenedione	29	<-2.0	<-2.0	M	M	S
5 $\beta$ -Androstenedione	30	<-2.0	<-2.0	M	M	S

5 Fig. 6 shows the canonical variate function for the pattern recognition obtained with the  $^{13}\text{C}$  NMR data only. The positive (upward-pointing) peaks in Fig. 6 correspond to bins that bias toward a strong RBA for binding to the estrogen receptor and negative (downwardly-pointing) peaks correspond to bins that bias toward a medium RBA. In Fig. 6, m/z bins 550 to 770 refer to the  $^{13}\text{C}$  NMR data from 0 ppm to 1 ppm for bin 550 to data from 221 ppm to 222 ppm for bin 770. The canonical variate function reveals that the aliphatic  $\text{CH}_2$  bins 580 to 585 (30 to 35 ppm) have a bias toward medium RBA. It also reveals that the

10

methyl CH<sub>3</sub> bins 558 and 566 (8 and 16 ppm, respectively) have a bias toward strong RBA. Many of the benzyl bins 665 to 700 (115 to 150 ppm) have a bias towards strong RBA. Thus these methyl groups and benzyl atoms are indicated to be important for strong RBA.

5           Based on the composite <sup>13</sup>C NMR and EI MS data, the statistical pattern-recognition program with 14 principal components (PCs) included 89.8% of the total variance and had a cross validation of 83.3%. The mathematical mode of significance was 97.6% at 1, with 28 degrees of freedom. The Wilks discriminant criterion was 99.89%. Fig. 7 shows the discriminant function for the composite  
10   <sup>13</sup>C NMR and EI MS data. Compounds with a white background are strong RBAs and compounds with a gray background are medium RBAs. Fig. 7 shows a large separation between 15 strong (left) and 9 medium (right) binders. There are 2 strong and 4 medium binders in the transition zone between the strong and medium binders. Table 1 shows that 25 of the 30 compounds are correctly predicted using  
15 the composite <sup>13</sup>C NMR and EI/MS data. Only 3β-androstanediol and hexestrol are incorrectly predicted using the canonical variate function. Again, the strong binding 3β-androstanediol is most likely predicted incorrectly because it is similar to the medium binding 3α-androstanediol.

          Fig. 8(a) shows the canonical variate function used in the pattern  
20 recognition of the EI MS data. Fig. 8(b) shows the canonical variate function used in the pattern recognition of the <sup>13</sup>C NMR data. The negative peaks in Figs 8(a) and 8(b) correspond to bins that bias toward a strong relative binder and positive peaks correspond to bins that bias toward a medium relative binder, to the estrogen receptor. In Figs. 8(a) and 8(b), the label "m/z bins 50 to 550" refers to the EI  
25 MS data, whereas "m/z bins 550 to 770" refers to the <sup>13</sup>C NMR data. In Fig. 8(a), the mass canonical variate is split evenly into bins that bias strong and medium binding. Many of the canonical variate bins that showed bias in Fig. 2 are also present in Fig. 8(b), but they are pointing in the opposite direction. The opposite directions found in Fig. 6 and 8(b) are insignificant technically, arising from the  
30 pattern-recognition program's arbitrary choice of left and right in the corresponding canonical variate score plot.

Only  $3\beta$ -androstanediol and hexestrol are incorrectly predicted in both SDAR models.  $3\beta$ -androstanediol is closest to  $3\alpha$ -androstanediol,  $5\alpha$ -androstanedione, and  $5\beta$ -androstanedione, all three of which have medium RBAs. Note however,  $3\beta$ -androstanediol has been inconsistently identified in the literature  
5 as a medium estrogen-receptor binder and a strong estrogen-receptor binder (see Miksicek, *J. Steroid. Biochem. Mol. Biol.* **49**:153-160, 1994 and Kuiper et al., *Endocrinology* **138**:863-870, 1997, respectively).

This example illustrates that, using the present method, compounds can be classified by biological activity based on similarities in the spectral patterns of the  
10 test compounds to the spectral patterns of the training set. In particular, additional test compounds can be classified according to their expected biological activity by obtaining the corresponding spectral pattern(s) of the test compound (e.g. NMR and MS patterns). Expected biological activity of the test compound can be predicted by detecting similarities between the spectral pattern of the test  
15 compound and the spectral patterns of compounds in the training set that are associated with a known biological activity. Similarities in patterns of the canonical variates associated with an endpoint (e.g. a biological activity such as estrogen receptor binding) derived for the training compounds with patterns in the test compound's segmented spectral data, are then used to predict whether the test  
20 compound will share the endpoint.

### Example 2 – Expanded Training Set for Estrogen Receptor Binding SDAR

An expanded training set of 119 compounds of varying estrogen receptor  
25 binding affinities was utilized to create an SDAR model from  $^{13}\text{C}$  NMR spectral data and to create an SDAR model from a composite of  $^{13}\text{C}$  NMR spectral data and EI MS spectral data. The training set included weak estrogen receptor binders in addition to strong and medium estrogen receptor binders. For this model, strong estrogen receptor binders were defined as having  $\log(\text{RBA}) > -0.25$  and weak  
30 estrogen receptor binders were defined as having  $\log(\text{RBA}) < -3.0$ . Medium estrogen receptor binders were defined as having  $\log(\text{RBA})$  in between. The

compound names, the input class ("Class"), the predicted endpoint class based upon  $^{13}\text{C}$  NMR only ("Class predicted NMR"), and the predicted endpoint class based upon  $^{13}\text{C}$  NMR plus EI MS ("Class predicted NMR/MS") for the training set are listed below in Table 2. Except for the increased number of compounds and a change in the mass spectral range utilized, the analysis was performed in the same manner as the analysis in Example 1.

Table 2

Compound	Log(RBA)	Class	Class predicted NMR	Class predicted NMR/MS
1,4-Diphenyl-1,3-butadiene		W	W	W
1,6-Dimethylnaphthalene		W	W	W
1,8-Octanediol		W	M	M
2,2'-Dihydroxy-4-methoxybenzophenone		W	W	W
2,2'-Methylenebis (4-chlorophenol)	-2.45	M	M	M
2,4,5-T		W	W	W
2,4-D		W	W	W
2,4-Dihydroxybenzophenone	-2.61	W	M	M
2-Chloro-4-methylphenol	-3.66	W	W	W
2-Chlorophenol	-3.67	W	W	W
2-Furaldehyde		W	W	W
2-OH-Estradiol *		S	S	S
2-Phenylphenol		W	W	W
2-sec-Butylphenol	-3.54	W	W	M
3-Phenylphenol	-3.44	W	W	W
4,4'-Dihydroxybenzophenone	-2.46	M	W	M
4,4'-Methylenebis(N,N-dimethyl)		W	M	W
4,4'-Methylenedianiline		W	W	W
4,4'-Methylene(2,6-ditertbutylphenol)		W	W	W
4,4'-Sulfonylphenol	-3.07	W	W	W
4-Benzyloxyphenol	-3.44	W	W	W
4-Chloro-2-methylphenol	-3.67	W	W	W
4-Chloro-3-methylphenol	-3.38	W	W	W



4-Ethylphenol	-4.17	W	W	W
4-Nonylphenol	-1.45	M	M	M
4-OH-Estradiol *		S	S	S
4-OH-Tamoxifen	2.24	S	S	W
4-Phenylphenol	-3.04	W	W	W
4-Stilbenol		W	W	S
4-tert-Amylphenol	-3.26	W	W	W
4-tert-Butylphenol	-3.61	W	W	W
4-tert-Octylphenol	-1.82	M	W	W
5 $\alpha$ -Androstane-3 $\alpha$ ,17 $\beta$ -diol	-2.67	M	W	M
5 $\alpha$ -Androstane-3 $\beta$ ,17 $\beta$ -diol	-0.92	M	M	M
Aldrin		W	W	W
Apigenin	-1.61	M	M	M
Aurin	-1.49	M	M	W
Benzylalcohol		W	W	W
Benzylbutyl phthalate		W	W	W
BiochaninA	-2.37	M	M	M
BIS (2-ethylhexyl) adipate		W	M	M
BIS (2-ethylhexyl) phthalate		W	W	W
BIS (2-hydroxyphenyl)-methane		W	W	W
BIS (4-hydroxyphenyl)-methane	-3.02	W	W	W
Bisphenol A	-2.11	M	M	M
Bisphenol B	-1.07	M	S	M
Butyl-4-aminobenzoate		W	W	W
n-Butylbenzene		W	W	W
Caffeine		W	M	W
Catechin		W	S	S
Chalcone	-2.82	W	S	W
Cholesterol		W	W	W
Chrysene		W	W	W
Chrysin		W	M	M
Cineole		W	W	W
Cinnamic acid		W	W	W
Clomiphene	-0.14	S	W	W
Corticosterone		W	W	S
Coumestrol	-0.05	S	S	S

DDD-o,p'		W	W	W
DDD-p,p'		W	W	W
DDE-p,p'		W	S	S
DDT-o,p'	-2.85	W	W	W
DDT-p,p'		W	W	W
DES	2.60	S	S	S
Daidzein	-1.65	M	M	M
Dexamethasone		W	S	W
Dibenzo-18-crown-6		W	W	W
Dibutyl phthalate		W	W	W
Dieldrin		W	S	S
Dienestrol	1.57	S	M	S
Diethyl phthalate		W	W	W
Dihydrotestosterone		W	M	W
Diisobutyl phthalate		W	W	S
Dimethyl phthalate		W	W	W
Diphenolic acid	-3.13	W	M	S
Dopamine		W	W	W
Estra-1,3,5(10)-trien-3-ol	1.14	S	S	W
Estra-1,3,5(10)-trien-3,6 $\alpha$ ,17 $\beta$ -triol	-0.15	S	S	S
Estriol	0.99	S	S	S
Estrone	0.86	S	S	S
Ethylcinnamate		W	W	W
Ethynyl estradiol	2.28	S	S	S
Etiocholan-17 $\beta$ -ol-3-one		W	S	S
Eugenol		W	W	W
Flavanone		W	W	M
Flavone		W	W	W
Genistein	-0.35	M	M	M
Heptanal		W	W	W
Hesperetin		W	M	M
Hexachlorobenzene		W	W	W
Hexestrol	2.48	S	W	W
Hexylalcohol		W	W	W
ICI-164,384	1.16	S	S	S
Isoeugenol		W	W	W

Kaempferol	-1.55	M	M	M
Lindane		W	W	S
Melatonin		W	M	W
Mestranol	0.35	S	S	S
Methoxychlor	-3.2	W	W	W
Morin	-3.35	W	W	W
Moxestrol	1.14	S	S	S
Nafoxidine	-0.14	S	S	S
Naringenin	-2.13	M	W	W
Naringin		W	W	W
Norethynodrel	-0.65	M	M	M
Phenolphthalein	-1.87	M	W	W
Phenol red	-3.25	W	W	W
Phloretin	-1.24	M	M	M
Progesterone		W	W	W
Quercetin		W	W	M
Suberic acid		W	W	W
Tamoxifen	0.21	S	S	S
Testosterone		W	W	W
Triphenylethylene	-2.78	W	W	W
Triphenylphosphate		W	W	W
Vanillin		W	W	W
17 $\alpha$ -Estradiol	0.49	S	S	S
17 $\beta$ -Estradiol	2.0	S	S	S

Based only on  $^{13}\text{C}$  NMR spectroscopic data, the statistical pattern recognition program with 22 principal components (PCs) had 88.9% of the total variance and a cross validation of 77.3 %. Figure 9 shows the discriminant function for  $^{13}\text{C}$  NMR data of 119 compounds. Compounds that are represented by an S are strong estrogen receptor binders, compounds that are represented by an M are medium estrogen receptor binders, and compounds that are represented by a W are weak estrogen receptor binders. Figure 9 shows a clustering between the 20 strong, 19 medium, and 80 weak binders. There are cluster overlaps between the weak and strong binders and between the weak and medium cluster

region. There are a total of eight false positives and two false negatives when using LOO to predict from the 119 compound SDAR model of estrogen receptor binding using only  $^{13}\text{C}$  NMR data.

Figure 10 shows the factors associated with the first canonical variate function for the pattern recognition of the  $^{13}\text{C}$  NMR data. The positive peaks in Figure 10 correspond to bins that bias toward a strong RBA for binding to the estrogen receptor and negative peaks correspond to bins that bias toward a medium RBA. In Figure 10,  $m/z$  bins 550 to 770 refer to the  $^{13}\text{C}$  NMR data from 0 ppm for bin 550 to 220 ppm for bin 770. The aliphatic  $\text{CH}_2$  bins 580 to 585 (30 to 35 ppm) have a bias toward medium RBA. The methyl  $\text{CH}_3$  bins 558 and 566 (8 and 16 ppm, respectively) have a bias toward strong RBA. Many of the aromatic bins 665 to 700 (115 to 150 ppm) have a bias towards strong RBA.

Based on the composite  $^{13}\text{C}$  NMR and EI/MS data over the range  $m/z = 100$  to  $m/z = 549$ , the statistical pattern recognition program with 22 principal components (PCs) included 80.1 % of the total variance and had a cross validation of 79.0 %. Figure 11 shows the discriminant function for  $^{13}\text{C}$  NMR data for 119 compounds. The compounds that are shown by an S exhibit strong RBAs, compounds that are shown by an M are exhibit medium RBAs, and compounds that are shown by a W exhibit weak RBAs. Figure 11 shows a similar clustering between the 20 strong, 19 medium, and 80 weak binders as seen Figure 9. There are cluster overlaps between the weak and strong binders and between the weak and medium cluster region. There are a total of nine false positives and three false negatives when using LOO to predict a compound's endpoint class from the 119 compound SDAR model of estrogen receptor binding using  $^{13}\text{C}$  NMR and EI/MS data.

Figures 12(a) and 12(b) shows the factors associated with the first canonical variate function for the pattern recognition of the composite  $^{13}\text{C}$  NMR and EI/MS data. The positive peaks in Figures 12(a) and 12(b) correspond to bins that bias toward a strong RBA for binding to the estrogen receptor and negative peaks correspond to bins that bias toward a medium RBA. In figure 12(a),  $m/z$  bins 100 to 549 are the EI/MS data bins and in Figure 12(b),  $m/z$  bins 550 to 770 refer to

the  $^{13}\text{C}$  NMR data from 0 ppm for bin 550 to 220 ppm for bin 770. The aliphatic  $\text{CH}_2$  bins 580 to 585 (30 to 35 ppm) have a bias toward medium RBA. The methyl  $\text{CH}_3$  bins 558 and 566 (8 and 16 ppm, respectively) have a bias toward strong RBA. Many of the aromatic bins 665 to 700 (115 to 150 ppm) have a bias towards  
5 strong RBA.

In both 119 compound SDAR models of estrogen receptor binding, a possible cutoff for testing experimentally whether a molecule did bind to the estrogen receptor would be anything with a positive component 1 as depicted in figures 9 and 11. This would leave no positives untested in each model. In both  
10 models, we would test 11 false negatives and cut the number of molecules to be tested for binding to the estrogen receptor to 31 out of the original 119 compounds.

A particularly striking result of the analysis using strong, medium, and weak estrogen receptor binders in the training set is the good separation between  
15 strong estrogen receptor binders and the other compounds as seen in Figures 9 and 11. There is a marked intermixing of the medium and weak estrogen receptor binders in Figures 9 and 11 with respect to the first canonical variate function (X-axis). These results are understandable in light of the physical basis for estrogen receptor binding. Estrogen receptor binding, like most ligand-receptor binding  
20 interactions, is governed by specific intermolecular interactions between the ligand and the receptor. The specific intermolecular interactions are what distinguish strong binders from the other compounds. Conversely, there are few or no specific interactions with the receptor on which to distinguish the medium and weak estrogen receptor binders from each other.

**Example 3 - Use of  $^{13}\text{C}$  NMR, EI MS, and Infrared Spectral Data to  
Produce a Predictive Model of Biodegradability for Monocyclic Chlorobenzene  
Derivatives**

5

Two recognized factors in the biodegradation of xenobiotic compounds in the environment are the physicochemical properties of the compounds themselves, and the enzymatic activities expressed by the microbial communities in the environment. The vast majority of microbial enzymatic reactions may be

10 classified into one of four categories: 1) group transfer; 2) oxidation and reduction; 3) elimination, isomerization, and rearrangement; and 4) carbon-carbon bond cleavage. The extent to which xenobiotic chemicals can serve as substrates for these enzymatic reactions is determined to a large extent by their structure and particularly by the presence of bonds that are similar to those found in the natural

15 substrates for the enzymes.

Infrared (IR) spectroscopic data may be included in an SDAR to provide a set of spectrally derived structure descriptors that include the types of bonds present in the subject molecules. Infrared data also reflect the modes and frequencies of vibration that are available to subject molecules. As chemical

20 reactions are often tied to particular vibrations of particular bonds, inclusion of the infrared data may improve the ability of an SDAR to predict reactivity.

The biodegradation data for many monocyclic chlorobenzene derivatives in sediment may be found in the Database for Environmental Fate of Chemicals ([www.aist.go.jp/RIODB/dbefc](http://www.aist.go.jp/RIODB/dbefc)). Additional data on the biodegradability is

25 published in *Biodegradation and Bioaccumulation Data of Existing Chemicals Based on the CSCL*, Japan Chemical Industry Ecology-Toxicology & Information Center (JETOC), Tokyo, Japan, 1992. The half-life period is used as the endpoint for the establishment of the SDAR and compounds are classified into two endpoint classes as readily biodegradable (R) (half-life < 30 days) and not readily

30 biodegradable (NR) (half-life > 30 days). The endpoint data for 34 chlorobenzene derivatives are given in Table 3 below.

Table 3

Compound	Half-life (days)	Input Class
Monochlorobenzene	46.2	NR
1,2-dichlorobenzene	36.9	NR
1,3-dichlorobenzene	433	NR
1,4-dichlorobenzene	385	NR
1,2,3-trichlorobenzene	23	R
1,2,4-trichlorobenzene	40.8	NR
1,3,5-trichlorobenzene	35	NR
1,2,3,4,-tetrachlorobenzene	18.2	R
1,2,3,5-tetrachlorobenzene	18.6	R
1,2,4,5-tetrachlorobenzene	28.8	R
Pentachlorobenzene	17.8	R
Hexachlorobenzene	27.1	R
2-chloroaniline	175	NR
3-chloroaniline	672	NR
4-chloroaniline	203	NR
2,3-dichloroaniline	54	NR
3,4-dichloroaniline	55	NR
3,5-dichloroaniline	149	NR
3,4,5-trichloroaniline	145	NR
2,3,4,5-trichloroaniline	57	NR
Pentachloroaniline	40	NR
2-chlorophenol	6.9	R
3-chlorophenol	29.2	R
4-chlorophenol	11.7	R
2,3-dichlorophenol	13.2	R
2,4-dichlorophenol	15.3	R
2,5-dichlorophenol	46.3	NR
2,6-dichlorophenol	5.0	R

3,4-dichlorophenol	69.9	NR
3,5-dichlorophenol	17.3	R
2,3,4-trichlorophenol	1.8	R
3,4,5-trichlorophenol	21.7	R
2,3,4,5-tetrachlorophenol	6.5	R
Pentachlorophenol	2.1	R

Spectral data for these chlorobenzene compounds is obtained from the Integrated Spectral Data Base System for Organic Compounds web site, [www.aist.go.jp/RIODB/SDBS](http://www.aist.go.jp/RIODB/SDBS), the *Aldrich Library of  $^{13}\text{C}$  and  $^1\text{H}$  FT NMR Spectra* (Poucher and Behnke, Eds., Aldrich Chemical Company, Volumes 1-3, 1993) and the NIST MS database software version 1.6. Experimental  $^{13}\text{C}$  NMR, EI MS, and IR data is collected when spectral data is not available in a database. Experimental spectral data is collected using standard spectroscopic protocols.

The spectral data points are segmented and used in the same way as 3D-QSAR uses comprehensive descriptors for structural and statistical analyses (CODESSA) (See, Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference). Specifically, the  $^{13}\text{C}$  NMR, EI-mass spectral, and IR data are placed into bins. Mass spectrometric data from m/z of 50 to 549 are used. Unassigned 1D  $^{13}\text{C}$  NMR chemical shifts are segmented into bins over a 0 to 221 ppm range. The  $^{13}\text{C}$  NMR frequencies are shifted to bins 550 to 770, so bin 550 is the  $^{13}\text{C}$  NMR spectrum for frequencies inside 0 to 1 ppm and bin 770 is the  $^{13}\text{C}$  NMR spectrum for frequencies inside 220 to 221 ppm. The unassigned IR frequencies in the range from  $4000\text{ cm}^{-1}$  to  $600\text{ cm}^{-1}$  are segmented into bins of  $50\text{ cm}^{-1}$  width. The IR data is shifted to bins 771 through 839 so that bin 771 holds the spectral data for the range  $4000\text{ cm}^{-1}$  to  $3950\text{ cm}^{-1}$  and bin 839 holds the spectral data for the range  $650\text{ cm}^{-1}$  to  $600\text{ cm}^{-1}$ . As an alternative, the IR data may be segmented into ranges of frequency or wavelength rather than the customary wavenumber frequencies.



To save space, the  $^{13}\text{C}$  NMR spectra are saved as sets of ordered pairs, each consisting of the respective chemical shift frequency in ppm and the number of peaks within the frequency range. Likewise, the IR spectra are saved as ordered pairs of bin number and the number of distinct IR peaks appearing in that particular range. Frequency ranges with only one peak are first assigned an area of 1; doubly degenerate frequencies an area of 2; and so forth. This is done to provide all the spectra with similar signal-to-noise ratios and to de-emphasize line-width variations due to differences in experimental conditions. The bin thus defines the number of significant and distinct spectral features within a frequency range. Normalization of the  $^{13}\text{C}$  NMR spectroscopic data and the IR data is accomplished by multiplying each set by a number that makes the maximum value in their respective set of bins equal to 100, which is the maximum value for EI MS data. The number of bins used to input  $^{13}\text{C}$  NMR spectra and IR spectra may be varied to improve the SDAR. Increasing the number of bins and shrinking the frequency width of each bin provides separate bins for closely spaced spectral features that may prove important as structure descriptors for establishing a reliable SDAR.

The segmented  $^{13}\text{C}$  NMR, EI MS, and IR spectral data along with the biodegradability data is input into the pattern recognition program RESolve Version 1.2 (Colorado School of Mines, Boulder, CO) and is auto-scaled and Fisher-weighted prior to principal component analysis (PCA). The discriminant analysis is performed based upon the canonical variate vectors. Leave-one-out (LOO) cross-validation is used to maximize the size of the training set. Spectral data for a test compound is then input and the compound judged to be readily biodegradable or not readily biodegradable based upon the presence of a majority of spectral features falling in bins that correspond to canonical variate factors from the training set that bias toward readily biodegradable or not readily biodegradable.

Alternatively, an expert system (self-learning or not) or artificial neural network may be utilized to perform the pattern recognition. Where the endpoint may be the result of multiple underlying mechanisms, such as multiple pathways of

biodegradation, expert systems or artificial neural networks may be advantageously utilized to separate compounds based upon their mode of biodegradation in addition to their biodegradability.

One example of an expert system is the MuRES sytem that is also part of the RESolve 1.2 software package. The MuRES package uses spectral data that is first compressed by projection onto a set of principal components. This method of compression is utilitarian, because it reduces the number of variables while maximizing the information content. MuRES uses the spectral scores that are calculated by projecting the spectral data onto a set of eigenvectors. However, MuRES may be applied directly without the compression step to data that already is overdetermined (i.e. more compounds and endpoint values than spectral data structure descriptors).

The knowledge base created by MuRES is in the form of simple binary rules. Binary rules use binary logic, logic that can be only true or false. A complex solution to a problem may be decomposed into a tree, often referred to as a classification tree, which consist of simple binary rules. The largest advantage of the expert system is that the scores do not have to be linearly separable which is an assumption required by discriminant analysis. Further details of the MuRES method are found in Harrington, *RESolve Software Manual*, Colorado School of Mines, Golden CO, which is incorporated herein by reference.

**Example 4 – Use of  $^{13}\text{C}$  NMR and UV/Visible Spectroscopic Data to Produce a Predictive Model for the Photosensitized Production of Singlet Molecular Oxygen.**

Phototosensitized oxidations involving singlet oxygen, a strong oxidant, are implicated in photodynamic inactivation of viruses and cells, in phototherapy for cancer, in photocarcinogenesis and in photodegradation of dyes and polymers. Quenching of excited singlet and triplet states of many substances by ground state molecular oxygen produces singlet oxygen; the lowest electronically excited singlet state of molecular oxygen. A compilation of the quantum yields for the formation of singlet oxygen in fluid solutions for over 700 substances is available from the

Notre Dame Radiation Laboratory – Radiation Chemistry Data Center  
(<http://www.rcdc.nd.edu>).

Compounds that are capable of photosensitizing the production of singlet oxygen are quite diverse and include aromatic hydrocarbons, aromatic ketones and thiones, quinones, coumarins, fluoresceins, transition metal complexes, and heterocyclics. Porphyrins and phthalocyanines are particularly important classes of compounds that are capable of producing singlet oxygen upon illumination.

To establish an SDAR useful for predicting the ability of molecules to photosensitize the production of singlet oxygen, the quantum yields for production of singlet oxygen by 20 compounds of diverse structure are obtained from the Radiation Chemistry Data Center website. The compounds are divided into two endpoint classes based upon having a high quantum yield for production of singlet oxygen (H, QY > 0.50) or a low quantum yield for production of singlet oxygen (L, QY < 0.50). Spectral data for these compounds is obtained from the resources mentioned in Examples 1 and 2 or is gleaned from other literature sources or is measured experimentally. The compounds, their singlet oxygen quantum yields, and the endpoint class for each are listed in Table 4.

Table 4

Compound	Quantum Yield	Endpoint Class
Acetophenone	0.29	L
Acridine	0.83	H
Anthracene	0.83	H
9,10-Anthraquinone	0.15	L
Anthrone	0.25	L
Azulene	0	L
Benzil	0.57	H
Benzophenone	0.29	L
Biacetyl	0.57	H

Biphenyl	1.0	H
Flourene	0.74	H
1-Methylindole	0.35	L
1,3-Diphenylisobenzofuran	0.26	L
Phenalen-1-one	0.94	H
Pivalothiophenone	1.0	H
Psoralen	0.0055	L
Pyrene	0.71	H
(E)-Stilbene	0.08	L
Tetracene	0.70	H
Xanthen-9-one	0.33	L

Ultraviolet-visible (UV-Vis) spectral data is used along with  $^{13}\text{C}$  NMR spectral data in the SDAR to include a measure of the importance of molecular excited states to the production of singlet oxygen. The segmented spectral data is used as a set of descriptors in the same way that 3D-QSAR uses comprehensive descriptors for structural and statistical analyses. (CODESSA) (See , Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein by reference). The  $^{13}\text{C}$  NMR spectral data is segmented into 1ppm bins over a 0 to 222ppm range. The UV-Vis data is segmented into 5 nm bins over the range of 190 nm to 900 nm. The  $^{13}\text{C}$  NMR spectral data occupies bins 1 through 221 and the UV-Vis data occupies bins 222 through 321. To save space, the  $^{13}\text{C}$  NMR spectral data is saved as sets of ordered pairs, each consisting of the bin number and the number of peaks with the frequency range corresponding to the bin. As described above, the  $^{13}\text{C}$  NMR bins define the number of significant and distinct spectral features within a frequency range. The UV-Vis data is saved as sets of ordered pairs, each consisting of the bin number and the average molar absorptivity of the molecule within the wavelength range corresponding to the bin. Molar absorptivity is used rather than absorbance to correct for variations in concentration between the measured UV-Vis spectra of the compounds.

The segmented  $^{13}\text{C}$  NMR and UV-Vis spectral data along with the quantum yields for singlet oxygen production for 20 compounds (the training set) are input as text files into the pattern recognition program RESolve Version 1.2 (Colorado School of Mines, Boulder CO). The spectroscopic data is auto-scaled and Fisher-weighted prior to principal component analysis (PCA). The discriminant analysis is performed based upon the canonical variate vectors. Leave-one-out (LOO) cross-validation is used to maximize the size of the training set and measure the ability of the SDAR to classify compounds correctly into their endpoint class. The number of bins used to input  $^{13}\text{C}$  NMR spectral data and UV-Vis spectral data may be varied. Increasing the number of bins will shrink the frequency width of each bin and provide separate bins for closely spaced spectral features that may prove important as structure descriptors for establishing a reliable SDAR. Increasing the number of compounds in the training set may also lead to a more reliable SDAR.

Once the training set has been cross-validated and deemed reliable, test compounds are then subjected to  $^{13}\text{C}$ -NMR and UV/Visible Spectroscopy, and these spectra are segmented into bins in the same manner as with the training set. Spectral patterns in these spectra for the test compound are then compared to the patterns for the training set, and endpoints associated with the spectral patterns of the training set are used to predict endpoints for the test compound. Where the endpoint is photosensitization of singlet oxygen, the presence or absence of spectral patterns in the training set which are associated with that endpoint are then detected to predict whether the test compound would be likely to have that characteristic.

#### **Example 5 – Use of an Experimentally Derived SDAR and $^{13}\text{C}$ NMR Spectral Data to Screen a Combinatorial Library of Compounds for Estrogen Receptor Binding Ability**

Combinatorial chemistry refers to methods of generating large numbers of compounds from smaller building block compounds that play an important role in generating lead compounds for rational drug design. The building block compounds are allowed to react to yield new compounds, either by mixed

synthesis using all building blocks together at once or by sequential reaction of the building block compounds. The reactions can take place within the virtual environment of a computer or they can actually be carried out in a reactor system. The resulting set of compounds is termed a combinatorial library.

5           A combinatorial library of compounds is produced using a combinatorial chemistry software package such as Afferent Structure™ (Afferent Systems, San Francisco, CA). With this software package it is possible to begin with a molecule and perform virtual synthetic steps on that molecule and any intermediate compounds made therefrom to yield a large number of intermediate and final  
10   product structures.

For each of the structures generated using the combinatorial chemistry software, <sup>13</sup>C NMR spectra are predicted. The <sup>13</sup>C NMR spectra may be predicted by any known method. Examples of methods for predicting <sup>13</sup>C NMR spectra include the neural network methods described by Kvasnicka (Kvasnicka, V., *J. Math. Chem.*, **6**: 63-76, 1991) and the quantum mechanical calculations of Dios et al. (Dios et al., *Science* **260**:1491-1496, 1993). Software for predicting <sup>13</sup>C NMR spectra is also available from Advanced Chemistry Development, Toronto, Ontario, Canada (www.acdlabs.com) (ACD/CNMR Spectrum Generator).  
15

The predicted <sup>13</sup>C NMR spectra are segmented into bins as the experimental  
20   <sup>13</sup>C NMR spectra in Example 1 were segmented. The segmented, predicted spectral data is input into the pattern recognition program and the SDAR established for estrogen receptor binding based on <sup>13</sup>C NMR only, from Example 1, is used to classify the compounds in the combinatorial library as either strong or medium estrogen receptor binders. Those compounds predicted by the SDAR to  
25   be strong estrogen receptor binders may then be tested experimentally for ability to bind to estrogen receptors.

Combinatorial libraries offer a quick approach to generating large numbers of new compounds, yet screening those compounds for specific biological activities is difficult and time consuming. The methods of the present invention provide  
30   rapid methods of screening the products produced by combinatorial chemistry

methods so that time consuming assays are performed only on compounds likely to exhibit the desired endpoint properties.

While this embodiment utilized an SDAR generated using experimental spectral data, in another embodiment, calculated spectral data is used to establish the SDAR. An SDAR established with calculated spectral data may be desirable when screening compounds according to their calculated spectral data, as the errors in calculated peak position are advantageously similar in both the training set spectral data and test compound spectral data.

#### 10      **Example 6 - Screening of Fractions for Specific Activities using SDAR**

The SDAR methods of the present invention are useful for screening raw fractions of compounds derived either from natural sources or from chemical reaction mixtures (including experimental combinatorial libraries).

A biological source of potentially new compounds, such as a sponge, is homogenized and partitioned to provide an aqueous fraction and an organic fraction. Each of these fractions is then chromatographed by any known method to yield a larger number of fractions that contain one or more compounds. Within some of these fractions, a compound or compounds capable of binding to estrogen receptors may be present. SDAR is used to quickly screen the fractions for the presence of estrogen receptor binding compounds.

An unassigned  $^{13}\text{C}$  NMR spectrum is obtained for each fraction and the spectrum is segmented into bins of 1ppm width. The fraction spectral data is then used along with the SDAR based on  $^{13}\text{C}$  NMR data only, as in Example 1, to predict the fraction's ability to bind to estrogen receptors. Screening of fractions in this manner avoids the time-consuming and costly step of subjecting each fraction to a bioassay.

Those fractions identified as containing compounds having the potential to bind to estrogen receptors may then be subjected to further analysis to reveal the number and identity of the compounds within the fraction.

### Example 7- Endpoints

Endpoints for use with the SDAR methods encompass the full range of biological, chemical, and physical properties exhibited by molecules. The methods of the present invention can be used to assist in drug design, biological activity  
5 predictions, toxicological predictions, chemical reactivity predictions, and metabolic pathway predictions. An endpoint is any molecular property or activity that can be measured qualitatively or quantitatively. Endpoints may be expressed in absolute or relative terms.

Endpoints may be chosen to establish SDARs that can be used to predict  
10 the environmental fate and toxicity of compounds. The ability of compounds to penetrate membranes, bind to enzyme active sites, react with soil, air, or water constituents, bind to soil constituents, hydrolyze, oxidize, and be transported in the environment can be used, along with spectral data for those compounds, to produce useful SDARs.

15 Spectral data can be used in combination with non-specific measures of toxicity, mutagenicity, teratogenicity, and carcinogenicity to establish SDARs. One example of such a non-specific measure is the Ames test. DNA damage and repair tests, Phosphorous-32 postlabeling, mutation induction in transgenes are others. Yet others include transgenic mouse assays, including the p53 +/- deficient  
20 model, the Tg.AC model, the TgHras2 model, and the XPA deficient model. LD<sub>50</sub> and EC<sub>50</sub> may provide endpoints for SDAR methods as well. Alternatively, the ability of compounds to induce specific biological outcomes such as cellular changes can be chosen as the endpoint used to establish the SDAR. For example, relevant tissues may be examined for changes at the cellular level using  
25 morphological, histochemical, or functional criteria. As appropriate, attention may be directed to such changes as the dose-relationships for apoptosis, cell proliferation, liver foci of cellular alteration, or changes in intercellular communication.

An SDAR may be established based upon any measurable response elicited  
30 in animals, plants, and microbes upon exposure to a series of compounds. Examples include SDARs based upon antiviral and antimicrobial activity. The



ability of compounds to induce metabolic disorders such as alterations in sugar metabolism may provide a useful endpoint. Phytotoxicity and stimulation of plant growth and reproduction are other examples. Pesticidal activity is yet another example. Measures of anti-hypertensive activity, anti-pyretic activity, anti-  
5 depressant activity, and the like further illustrate useful endpoints that are usually related to human health.

Phototoxicity, both specific and non-specific, may be correlated with spectral features to yield an SDAR.

Multiple endpoints may be utilized to establish multiple SDARs from a  
10 single set of spectral data. Compounds then may be screened based upon their spectra using multiple SDARs for any combination of desirable or undesirable activities. One example of a useful combination is that of maximal potential efficacy as a therapeutic agent with minimal potential side effects. Agrochemicals may be screened using multiple SDARs for species-specific toxicities and  
15 tolerances.

An especially useful application of the methods of the present invention is to the prediction of ligand-target molecule binding. The binding of a molecule to a target such a protein, nucleic acid, synthetic polymer, chimeric molecule, or membrane constituent is often the most important step in the elicitation of a  
20 particular property or activity by a molecule. Binding affinities for ligand-target molecule interactions can be expressed in either absolute (e.g., an equilibrium constant) or in relative (e.g., relative to a reference compound, as determined for example by a competitive binding experiment) terms. Example 1 above is one example of how the relative binding affinity of a series of molecules can be utilized  
25 along with spectral data to establish a predictive SDAR model. SDAR models based upon relative binding affinities may be useful for rapidly and inexpensively screening compounds for a particular activity. They also may be useful tools for rational drug design when used to identify the spectral, and thus structural, features responsible for that activity.

30 The metabolic pathway involved in the production or destruction of a series of molecules is another endpoint useful for the methods of the present invention.

A predictive SDAR based upon pathway-structure relationships may be able to predict the biosynthetic path for newly discovered naturally occurring compounds. Similarly, SDAR using biodegradability as an endpoint may be useful for predicting the residence time of pollutants in the environment.

5 Rates of reaction and other measures of reactivity, such as site of reaction on a molecular structure, including the site of electrophilic aromatic substitutions on aromatic compounds, are useful chemical endpoints for the practice of the present invention.

Physical constants such as water-octanol partition coefficients, vapor  
10 pressures, pKa, pKb, hydrophobicities, relative acidities and basicities as well as water solubilities can be used with spectral data to provide SDARs. Such estimates may be especially useful for physical properties that are difficult and time-consuming to measure. For example, octanol-water partition coefficients are important for modeling the environmental transport of chemicals. While the  
15 octanol-water partition coefficient of a compound might be available, it is less likely that transient species derived from that compound during biodegradation are available in sufficient quantities to measure their octanol-water partition coefficients. SDAR according to the methods of the present invention provides an efficient way to predict the octanol-water partition coefficient for transient species,  
20 whose environmental transport characteristics need to be modeled.

Other examples of endpoints useful for the methods of the present invention, in addition to those discussed above, may be found in Hansch and Leo, *Exploring QSAR: Fundamentals and Applications in Chemistry and Biology*, American Chemical Society, 1995. Further examples of endpoints useful for the  
25 methods of the present invention may be found in *Quantitative Structure-Activity Relationships in Environmental Sciences-VII*, Chen and Schüürmann, eds., SETAC Press, 1997.

### Example 8 – Spectral Data

30 Spectroscopy refers to branch of analytical chemistry in which atomic and molecular structure is studied by measuring radiant energy absorbed or emitted by

a substance in any of the wavelengths of the electromagnetic spectrum, in response to excitation by an external energy source. The types of absorption and emission spectroscopy are usually identified by the wavelength involved, such as gamma-ray, X-ray, UV, visible, infrared, microwave, and radiofrequency. Nuclear magnetic resonance spectroscopy (NMR) examines differences in energy states created by a magnetic field. Spectral data refers to the measurements of the energy differences across the spectrum, and spectral patterns refer to differences in the detected energy differences measured across a region of the electromagnetic spectrum. Any instrumental method that produces data that depend upon the structural and quantum mechanical properties of a molecule may be utilized with the methods of the present invention.

Spectral data as used in some embodiments includes the entire spectrum (or spectra) generated by the instrumental method (or methods) of spectroscopy or by calculation. Furthermore, the spectral data need not be assigned to particular structural features. In other embodiments the spectral data comprises only a portion of the spectrum or spectra available. The spectral portions utilized in the methods of the present invention may conveniently cover a spectral region known to typically arise from one or more particular structural features. For example, with respect to  $^{13}\text{C}$  NMR spectral data, spectral data can be obtained from the entire  $^{13}\text{C}$  NMR spectrum (0 to 220 ppm), or at least half or a third of that spectrum, or at least a 60 ppm, 80 ppm, 100, or 150 ppm portion of the spectrum. With respect to IR spectral data, for example, the spectral data can be obtained from the entire IR spectrum ( $4000\text{ cm}^{-1}$  to  $500\text{ cm}^{-1}$ ), or at least a hundredth, fiftieth, quarter, or half of that spectrum, or at least a 35, 50, 100, 200, 500, or 1000  $\text{cm}^{-1}$  portion of the spectrum. In particular embodiments, all or substantially all of the spectral data within a particular portion of the spectrum is obtained. However, the spectral features within the entire spectrum or within the portion of the spectrum need not be assigned to structural features and referenced to the corresponding spectral features arising from the structure features of a reference compound. For example, rather than utilizing the assigned  $^{13}\text{C}$  NMR resonances of the carbon atoms in a substituted benzene ring referenced to the  $^{13}\text{C}$  NMR

resonances of the carbon atoms in unsubstituted benzene, the present invention could utilize the entire spectral region wherein benzene ring carbons typically fall, without assigning and referencing the data therein to the corresponding spectral data for benzene, which has a known structure.

5 Nuclear magnetic resonance (NMR) data is especially attractive for use with the present invention because of the large amount of structural information contained in a NMR spectrum. NMR instrumentation is widely available and NMR spectra are obtained routinely during structure elucidation. Additionally, the NMR spectra of many compounds have already been measured and are available  
10 (see Example 1 for some representative sources).

$^{13}\text{C}$  NMR and  $^1\text{H}$  NMR spectral data (especially  $^{13}\text{C}$  NMR spectral data) are very sensitive to subtle changes in substitution, conformation, chirality, and electronic density. Moreover, changes in  $^{13}\text{C}$  NMR chemical shifts can occur at a site as many as five carbon atoms removed from the site of the variation.  
15 Solvation and proton-exchange effects on the electronic properties of molecules are more clearly reflected in  $^1\text{H}$  NMR chemical shifts and line widths. One-dimensional  $^{13}\text{C}$  NMR and  $^1\text{H}$  NMR spectral data as well as two-dimensional  $^{13}\text{C}$ - $^1\text{H}$  heterocorrelation data such as that derived from HSQC and HMQC experiments are useful. Furthermore, modern FT-NMR instruments are capable  
20 of providing NMR spectral data for as little as 1 nanogram of a compound (an amount likely insufficient for performing standard bioassays such as the Ames test).

NMR data may be segmented into bins prior to their analysis, along with endpoint data, in a pattern-recognition program. Suitable bin widths will vary  
25 according to the identity of the nuclei for which the spectrum is generated, and whether the technique is one or two-dimensional. For one-dimensional  $^{13}\text{C}$  NMR spectral data, the bin width may be varied from the digital resolution of the instrument (typically about 0.1 ppm) to about 50ppm. For one-dimensional  $^1\text{H}$  NMR spectral data, the bin width may be varied from the instrumental digital  
30 resolution (typically about 0.01 ppm) to about 2 ppm. For two-dimensional  $^{13}\text{C}$ - $^1\text{H}$  heterocorrelation data, the bin may be defined by similar corresponding widths in

both the  $^{13}\text{C}$  and  $^1\text{H}$  dimensions. Even data of higher dimensions (e.g., three, four, etc.) including NMR spectral data from other nuclei, such as  $^{15}\text{N}$ ,  $^{31}\text{P}$ ,  $^{19}\text{F}$ ,  $^{17}\text{O}$ , and  $^{35}\text{S}$  may be used in establishing an SDAR. Correspondingly, bins may be defined with respect to each dimension and may be of a width equal to the digital  
5 resolution of the data or greater.

In another embodiment of the invention,  $^{13}\text{C}$  NMR spectral data are predicted by calculation (see, for example, Dios et al., *Science* **260**:1491-1496, 1993 and Kvasnicka, V., *J. Math. Chem.*, **6**: 63-76, 1991) and used in an SDAR model that has been trained on true  $^{13}\text{C}$  NMR spectral data. Software for  
10 predicting  $^{13}\text{C}$  NMR spectra is also available from Advanced Chemistry Development, Toronto, Ontario, Canada (www.acdlabs.com) (ACD/CNMR Spectrum Generator). Predicted  $^{13}\text{C}$  NMR spectral data may be used, for example, to aid in rational drug design by allowing proposed structures to be tested for potential activities before synthesis is attempted.

15 If predicted spectral data is utilized to establish an SDAR, the spectral data may be segmented into bins that are desirably of a width equal to the average standard deviation in chemical shift predicted by the method, or greater. Likewise, test structure predicted spectra may be segmented in a similar fashion.

Mass spectrometry can provide a measure of the size of a molecule, the  
20 size and identity of a molecule's structural subunits, and information regarding bond strengths within a molecule. Mass spectral data, especially electron impact mass spectral (EI MS) data, has already been obtained for many compounds and, even more so than NMR data, is available from convenient sources (see Example 1). EI MS data is also a standard technique used in structure elucidation.

25 Other mass spectrometric techniques that are useful for providing additional and often complementary information include time-of-flight mass spectrometry (TOF MS), chemical ionization mass spectrometry (CI MS), fast-atom bombardment (FAB). Modern TOF MS spectrometers are capable of providing mass-spectral data from 1 ng or less of purified material (an amount that is likely  
30 insufficient for performing standard activity screens such as the Ames test).

Mass spectral data may be segmented into ranges of  $m/z$  ratio (for instance, ranges corresponding to mass ranges from about the digital resolution of the instrumental method, typically about 0.1 amu, to about 50 amu) or may be segmented according to integer  $m/z$  ratios, with non-integer  $m/z$  ratios being  
5 rounded to the nearest integer.

Infrared (IR) spectra may also be used, for example in establishing SDAR models capable of discerning differences in activity seen for tautomers that are indistinguishable in NMR and mass spectrometric data. Infrared (IR) spectra are treated in a similar fashion to the NMR spectral data of Example 1 in that each  
10 spectrum may be separated into bins of a certain spectral range, for example from about  $1\text{ cm}^{-1}$  to about  $200\text{ cm}^{-1}$  for entry into a pattern-recognition program

Ultraviolet-Visible (UV-Vis) spectral data, which is inherently reflective of the electronic energy levels of a molecule, may be used by segmenting the spectral data into bins of a certain spectral range, for example from about the digital  
15 resolution of the instrumental method to about 50 nm. UV-Vis spectral data can be used, for example, in predicting phototoxicity under solar illumination. Similarly, fluorescence and phosphorescence spectra may be handled analogously to UV-Vis spectra and utilized to establish an SDAR. Fluorescence and phosphorescence spectra reflect the energy redistribution within a molecule upon  
20 absorption of light and thus may provide important structure descriptors for predicting the light driven properties of molecules.

In addition to the possibility of using a single type of spectral data to establish an SDAR, spectral data of various types may be combined to form composite sets of spectral data. Entire spectra or particular regions of spectra may  
25 be combined to yield spectral data sets that may be used in the methods of the present invention, along with endpoint data, to establish the SDAR. Spectral data may come from any composite of NMR, MS, IR, Fluorescence, Phosphorescence, and UV-Vis spectra, including composites of different species of spectra within these broad genera of spectra. Furthermore, different regions of any type of  
30 spectrum can be segmented into bins of different sizes so, for example, portions of a spectrum with many closely spaced peaks can be described by narrow spectral

bins and portions of a spectrum without many peaks can be described by wide spectral bins.

In some embodiments the spectral data is not used in its raw form to establish an SDAR, but rather the data is subjected to pattern recognition analysis after some sort of pre-treatment to improve the ability of pattern recognition to extract the SDAR. For example normalization may be used to equalize the importance of spectral data derived from different instrumental methods when forming a composite, such as, a composite of MS data and NMR data wherein the maximum signals might be 100 and 1000 respectively. Autoscaling may be used to equalize the importance of inherently weak spectral data with inherently strong spectral data, for example, UV-Vis absorption bands within an absorption spectrum with very different extinction coefficients. Fisher-weighting may be used to emphasize the spectral data that are most important for predicting the endpoint data, such as spectral data found in compounds with a large endpoint values but absent from compounds with small endpoint values.

In general, techniques for pre-treating data include artifact removal and/or linearization, centering, and scaling and weighting. A common form of artifact removal is baseline correction of a spectrum. Common linearizations include the conversion of spectral transmittance into spectral absorbance and the multiplicative scatter correction for diffuse reflectance spectra. Centering, sometimes called mean centering is simply the subtraction of the mean spectral signal at each frequency or  $m/z$  from each spectrum. Scaling or weighting involves multiplying all of the spectra by a different scaling factor for each sub-spectral region. This is done to increase or decrease the influence of certain spectral regions or features. A particular example of weighting is Fisher-weighting. Two types of scaling are typically encountered, variance scaling and autoscaling. Further discussion of data pretreatment may be found in Kramer, R., *Chemometric Techniques for Quantitative Analysis*, Marcel Dekker, Inc., 1998. Additional methods for the pre-treatment of data prior to pattern recognition are known in the art and are included in the present invention.

Since spectral data is utilized in some embodiments in the same manner as 3D-QSAR utilizes comprehensive descriptors for structural and statistical analyses (CODESSA) (see, Tong et al., *J. Med. Chem.*, **39**: 380-387, 1995 and Collantes et al., *J. Anal. Chem.*, **68**: 2038-2043, 1996, both of which are incorporated herein  
5 by reference) it is possible to combine the spectrally derived structure descriptors of the present invention with structure descriptors based on bulk measurements (such as the octanol/water partition coefficient) and still retain the ability to produce predictive models without the need for structural knowledge beforehand. Hybrid methods also fall within the scope of the invention wherein the spectrally  
10 derived descriptors, which do not require structural knowledge beforehand, may be combined with other structure descriptors that require structural knowledge to produce a larger set of descriptors for use in a predictive model.

### Example 9 – Pattern-Recognition Programs

15 Pattern-recognition programs useful for practicing the present invention are of two major types; statistical and artificial intelligence.

Statistical methods include Principal Component Analysis (PCA) and variations of PCA such as linear regression analysis, cluster analysis, canonical variates, and discriminant analysis, soft independent models of class analogy  
20 (SIMCA), expert systems, and auto spin (see, for example, Harrington, *RESolve Software Manual*, Colorado School of Mines, 1988, incorporated by reference). Other examples of statistical analysis software available for principal-component-based methods include SPSS (SPSS Inc., Chicago, IL), JMP (SAS Inc., Cary NC), Stata (Stata Inc., College Station, TX) and Cluster (available to run from  
25 entropy:~dblank/public\_html/cluster).

Artificial intelligence methods include neural networks and fuzzy logic. Neural networks may be one-layer or multilayer in architecture (See, for example, Zupan and Gasteiger, *Neural Networks for Chemists*, VCH, 1993, incorporated herein by reference). Examples of one-layer networks include Hopfield networks,  
30 Adaptive Bidirectional Associative Memory (ABAM), and Kohonen Networks. Examples of Multilayer Networks include those that learn by counter-propagation



and back-propagation of error. Artificial neural network software is available from, among other sources, Neurodimension, Inc., Gainesville, FL (Neurosolutions) and The Mathworks, Inc., Natick, MA (MATLAB Neural Network Toolbox).

- 5           Spectral patterns can be analyzed using other approaches. Analog spectral peak patterns may be digitized, and image analysis may be used to search for similarities or differences between the spectral patterns of training sets and test compounds

10           **Example 10 – Computer Environments for SDAR Methods.**

The SDAR methods of the present invention may be implemented using a single computer or utilizing a distributed computing environment.

*Exemplary Distributed Computing Environment*

- Fig. 13 illustrates a distributed computing environment in which the software elements used to implement the SDAR methods of the present invention may reside. The distributed computing environment 100 includes two computer systems 102, 104 connected by a connection medium 106. The computer systems 102, 104 can be any of several types of computer system configurations, including personal computers, multiprocessor systems, and the like. In terms of logical relation with other computer systems, a computer system can be a client, a server, a router, a peer device, or other common network node. Moreover, although Fig. 13 illustrates two computer systems 102, 104, the present invention is equally applicable to an arbitrary, larger number of computer systems connected by the connection medium 106. Additional computer systems 102 or 104 may be connected by an arbitrary number of connection mediums 106. The connection medium 106 can comprise any local area network (LAN), wide area network (WAN), or other computer network, including but not limited to Ethernets, enterprise-wide computer networks, intranets and the Internet.

- Portions of the SDAR software can be implemented in a single computer system 102 or 104, with the application later distributed to other computer systems 102, 104 in the distributed computing environment 100. Portions of the SDAR

software may also be practiced in a distributed computing environment 100 where tasks are performed by a single computer system 102 or 104 acting as a remote processing device that is accessed through a communications network, with the distributed application later distributed to other computer systems in the distributed computing environment 100. In a networked environment, program modules comprising the SDAR software can be located on more than one computer system 102 or 104. Communication between the computer systems in the distributed computing network may advantageously include encryption of the communicated data.

### *Exemplary Computer System*

Fig. 10 illustrates an example of a computer system 120 that can serve as an operating environment for the SDAR software. With reference to Fig. 14 an exemplary computer system for implementing the invention includes a computer 120 (such as a personal computer, laptop, palmtop, set-top, server, mainframe, and other varieties of computer), including a processing unit 121, a system memory 122, and a system bus 123 that couples various system components including the system memory to the processing unit 121. The processing unit can be any of various commercially available processors, including Intel x86, Pentium and compatible microprocessors from Intel and others, including Cyrix, AMD and Nexgen; Alpha from Digital; MIPS from MIPS Technology, NEC, IDT, Siemens, and others; and the PowerPC from IBM and Motorola. Dual microprocessors and other multi-processor architectures also can be used as the processing unit 121.

The system bus can be any of several types of bus structure including a memory bus or memory controller, a peripheral bus, and a local bus using any of a variety of conventional bus architectures such as PCI, VESA, AGP, Microchannel, ISA and EISA, to name a few. The system memory includes read only memory (ROM) 124 and random access memory (RAM) 125. A basic input/output system (BIOS), containing the basic routines that help to transfer information between elements within the computer 120, such as during start-up, is stored in ROM 124.

The computer 120 further includes a hard disk drive 127, a magnetic disk drive 128, e.g., to read from or write to a removable disk 129, and an optical disk drive 130, e.g., for reading a CD-ROM disk 131 or to read from or write to other optical media. The hard disk drive 127, magnetic disk drive 128, and optical disk drive 130 are connected to the system bus 123 by a hard disk drive interface 132, a magnetic disk drive interface 133, and an optical drive interface 134, respectively. The drives and their associated computer-readable media provide nonvolatile storage of data, data structures, computer-executable instructions, etc. for the computer 120. Although the description of computer-readable media above refers to a hard disk, a removable magnetic disk and a CD, it should be appreciated by those skilled in the art that other types of media which are readable by a computer, such as magnetic cassettes, flash memory cards, digital video disks, Bernoulli cartridges, and the like, can also be used in the exemplary operating environment.

A number of the SDAR program modules can be stored in the drives and RAM 125, including an operating system 135, one or more application programs 136, other program modules 137, and program data 138.

A user can enter commands and information into the computer 120 through a keyboard 140 and pointing device, such as a mouse 142. Other input devices (not shown) can include a microphone, satellite dish, scanner, or the like. These and other input devices are often connected to the processing unit 121 through a serial port interface 146 that is coupled to the system bus, but can be connected by other interfaces, such as a parallel port, game port or a universal serial bus (USB). A monitor 147 or other type of display device is also connected to the system bus 123 via an interface, such as a video adapter 148. In addition to the monitor, computers typically include other peripheral output devices (not shown), such as printers.

The computer 120 can operate in a networked environment using logical connections to one or more other computer systems, such as computer 102. The other computer systems can be servers, routers, peer devices or other common network nodes, and typically include many or all of the elements described relative to the computer 120, although only a memory storage device 149 has been

illustrated in Fig. 14. The logical connections depicted in Fig. 10 include a local area network (LAN) 151 and a wide area network (WAN) 152. Such networking environments are common in offices, enterprise-wide computer networks, intranets and the Internet.

5           When used in a LAN networking environment, the computer 120 is connected to the local network 151 through a network interface or adapter 153. When used in a WAN networking environment, the computer 120 typically includes a modem 154 or other means for establishing communications (e.g., via the LAN 151 and a gateway or proxy server 155) over the wide area network 152, 10 such as the Internet. The modem 154, which can be internal or external, is connected to the system bus 123 via the serial port interface 146. In a networked environment, program modules depicted relative to the computer 120, or portions thereof, can be stored in the remote memory storage device. It will be appreciated that the network connections shown are exemplary and other means of establishing 15 a communications link between the computer systems (including an Ethernet card, ISDN terminal adapter, ADSL modem, 10BaseT adapter, 100BaseT adapter, ATM adapter, or the like) can be used.

          In accordance with the practices of persons skilled in the art of computer programming, a particular embodiment of the SDAR method is described in Fig. 4 20 with reference to acts and symbolic representations of operations that may be performed by the computer 120. Such acts and operations are sometimes referred to as being computer-executed. It will be appreciated that the acts and symbolically represented operations include the manipulation by the processing unit 121 of electrical signals representing data bits which causes a resulting 25 transformation or reduction of the electrical signal representation, and the maintenance of data bits at memory locations in the memory system (including the system memory 122, hard drive 127, floppy disks 129, and CD-ROM 131) to thereby reconfigure or otherwise alter the computer system's operation, as well as other processing of signals. The memory locations where data bits are maintained 30 are physical locations that have particular electrical, magnetic, or optical properties corresponding to the data bits.

In view of the many possible embodiments to which the principles of our invention may be applied, it should be recognized that the illustrated embodiments are only specific examples of the invention and should not be taken as a limitation on the scope of the invention. Rather, the scope of the invention is defined by the  
5 following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2